THE AFRICAN SPECIES OF CULEX AND ALLIED GENERA. By F. W. Edwards, B.A.

(Published by permission of the Trustees of the British Museum.)

The writer's original intention in publishing this paper was to give a key for the identification of those African mosquitos which might be taken by the nonspecialist for species of *Culex*, without attempting to form any opinions as to the value or limits of allied genera. A study of the insects, however, led to the realisation that scale characters, in this group at least, are not of very great value for separating genera, and consequently the present paper gives the author's views of the classification of the *Culex* group, and is more complete than it might otherwise have been. Time has not yet been available to deal with those species which have flat scales on the top of the head or on the sentellum (*Stegomyia* group).

It will be noticed at once that a large number of names (nearly seventy) have been sunk as synonyms, and it may be as well to point out that this has, as a rule, only been done after a careful comparison of the types. Where there is room for doubt as to the synonymy of two names, this has been freely admitted.

All that is claimed for this paper is that it may make the determination of good specimens a little easier than up to the present. The group of species including *Culex invidiosus*, and some other species, are very difficult to classify, and the author has not been able, much to his regret, to find any clearly marked distinctions.

The CULICIDAE as a whole are classified as follows by Lt.-Col. Alcock (Ann. Mag. Nat. Hist. (8) viii., p. 241, 1911) :--

Sub-family A. Corethrinae. (Proboseis short.)

Sub-family B. Culicinae. (Proboscis long.)

Section 1. MEGALORHINI. (Proboscis very long and bent downwards. Usnally large metallic species.)

Section 2. EPIALURGI = Anophelinae of Theobald. (Palpi long in Q.)

- Section 3. CULICALES, comprising Culicinae, Heptaphlebomyinae, Dinoceratinae, Aedinae and Uranotaeniinae of Theobald. (Metanotum bare.)
- Section 4. METANOTOTRICHA, comprising Trichoprosoponinae. Dendromyinae and Limatinae of Theobald. (Metanotum bearing bristles.)*

This grouping of the main divisions of the family is in the present writer's opinion the best that has yet been proposed. Many recent authors have adopted an essentially similar plan, but have made only three divisions of the CULICINAE (or, where the CORETHRINAE are regarded as a separate family, three sub-families of CULICIDAE), the MEGALORHINI (Megarhininae of Theobald) being included with the CULICALES. It seems better, however, owing to the many differences of structure between the two groups, to keep them separate.

^o [It has very reasonably been suggested to me that it is advisable to use similar terminations for the names given to groups of equal value; and further, that it is inadvisable to interfere with the established name of "Anopheline" for the malaria-carrying mosquitos. Mr. Guy Marshall proposes the names (1) Megalorhinina, (2) Anophelina, (3) Culicina, and (4) Metanototrichina, for the four sections of CULICINAE, and with this proposal 1 entirely concur.— A. Alcock.]

The CULICALES may be divided further into two main groups as follows :— Sub-section I. CULICIDES. Palpi of \mathcal{J} longer than those of \mathcal{Q} .

Sub-section II. AEDIDES. Palpi similar in both sexes.

Further sub-division, except into genera, is unnecessary, and probably not

In the following table are included all those forms of the CULICIDES (Theobald's sub-families *Culicinae* and *Heptaphlebomyinae*) which have no flat scales on the scutellum or on the middle part of the head. The genus *Ludlowia* has also been included owing to the similarity of the females to *Culex*, and the fact that not all the species have a flat-scaled head.

Table of Genera.

1.	Male palpi much shorter than proboseis	2.
	,, never much shorter than proboseis (in Theobaldia	
	spathipalpis they often shrink and curl up, so as to	
	appear shorter than the proboseis)	3.
2.	Ungues all simple in both sexes PROTOMELANOCONION, p.	244.
	Fore and mid ungnes dentate in both sexes [two undescribed spec	ies].
3.	Male palpi apparently 2-jointed	4.
	", ", 3-jointed	5.
4.	Fore and mid ungues of female simple; first fork-cell in female	
	shorter than second; wings nearly bare on lower half	
	LUDLOWIA, p.	244.
	Fore and mid nugnes of female dentate; first fork-cell in female	
_	longer than second; thorax deep yellow at sides BANKSINELLA, p.	245.
5.	Last two joints of male palpi thickened, terminal usually shorter	
	than penultimate and turned slightly downwards (fig. 1)	6.
2	Last two joints of male palpi thin, nucronate and curved npwards	9.
6.	Fore and mid ungues of male and female dentate (both claws)	7.
	All ungues of female, and hind pair of male, simple	8.
1.	remate palpi more than half as long as probosels; wing-scales	010
	broad, legs densely clothed with ontstanding scales MUCIDUS, p.	246.
	remale pappi less than half as long as probosels; wing-scales	040
U U	mostly narrow; legs without outstanding scales OCHLEROTATUS, p.	240.
с .	Turse species	951
	Smaller day on fore and mid taxi of mule not toothed a medium-	201
	sized usually vollow species TAENIOPHYNCHUS D	951
9	Terminal joint of male value very short (fig 2): wing-scales all	201.
47.0	broad MANSONIOLDES D	253
	Terminal joint of male nalpi nearly as long as or longer than	-00°.
	penultimate : wing-scales mostly narrow	10.
10	Male palpi with a row of ontstanding transparent whitish seales : a	
	row of flat on alescent scales round the eve-margins in both sexes	
	Culiciomyia, D.	254.
	Male palpi without such scales (fig. 3); flat scales not continuous	
	round eye-margins CULEX, p.	256.
	Totala of changelie the the the the of Lan, pr	

 $\mathbf{242}$

advisable.

Table of Females.

(Species of the Aedes group (not included in this table), though obscure-looking. nearly always have a flat-scaled head). 1. Fore and mid ungues toothed 2. All ungues simple 5. 2. Hind ungues toothed 3. ", " simple … … … … ... 4. . . . 3. Palpi more than half length of proboscis : wing scales broad ; legs shaggy Mucidus, p. 246. Palpi less than half length of proboscis; wing scales narrow; Ochlerotatus, p 246. legs not shaggy ... BANKSINELLA, p. 245. 4. Thorax bright yellow at the sides, not bright yellow at sides ... OCHLEROTATUS, p. 246. 5. Distinctly yellow species ; legs largely yellow TAENIORHYNCHUS, p. 251. Not yellow species 6. 6. Wing-seales all very broad (compare also Aedeomyia) MANSONIOIDES, p. 253. mostly narrow 7. 7. Head mostly covered with flat scales (as in Stegomyia, etc.) LUDLOWIA, p. 244. ", ", narrow eurved scales 8. 8. Head with a row of small flat seales round the eye-margins CULICIOMYIA, p. 254. Head without flat scales on the occipit 9. 9. Last two joints of hind tarsi white LUDLOWIA (MEGACULEX), p. 244. , , , not white 10. 10. Large to medium-sized species ; femora and tibiae with pale spots 11. Femora and tibiae not spotted 13. 11. Thorax with distinct narrow white lines; a pale ring before tips of femora Theo. spathipalpis, p. 251. Thorax without white lines 12. 12. Thorax with the anterior two-thirds, or at least the middle third, with pale seales C. quasigelidus, p. 258. Thorax not so marked ; spots on legs more numerous C. tigripes, p. 261. 13. Thorax with the anterior two-thirds bearing pale seales ... 14. " almost nniformly scaled 16. ... 14. Violet-blackish species *Taen. metallicus*, p. 252. Not violet-blackish species 15. 15. Abdominal segments with pale median basal spots C. annulioris, p. 259. ,, without such spots C. consimilis, p. 259. 16. A well-defined margin of white scales to eyes; small grey species. with small whitish basal spots on abdominal segments Prot. juscum, p. 244. No well-defined rim of white seales to eyes (except sometimes in C. grahami) 17. 17. Large species, joints of tarsi basally banded Theo. annulata, p. 251. Medinm-sized or small species, tarsi not banded, or rarely with basal and apical bands ... CULEX, p. 256.

Genus PROTOMELANOCONION, Theo.

Mon. Cul. V, p. 462 (1910).

1. P. fuscum, Theo., Mon. Cul. V, p. 463 (1910).

Although the generic characters will at once distinguish the male from all the related genera, the female is a very obscure insect, and might easily be mistaken for a small *Culex*, a *Culiciomyia*, or an *Aedes*. It perhaps bears the greatest resemblance to a *Culiciomyia*, since it has the same rim of white scales round the eye-margins : but an examination with the microscope shows that the white scales are narrow-curved and not flat. *P. fuscum* φ can be distinguished from *Culex grahami* and its allies by the much longer bristles on the thorax.

Genus LUDLOWIA, Theo.

Mon. Cnl. IV, p. 193 (1907).

Megaculex, Theo., Mon. Cul. IV, p. 282 (1907). Radioculex, Theo., Rec. Ind. Mus. 11, p. 295 (1908). Hispidimyia, Theo., Mon. Cul. V, p. 245 (1910).

The essential characters of the genus are :—(1) the apparently 2-jointed \circ palpi, of which the apical joint is swollen; (2) the short fork-cells, especially the first; (3) the rather peculiar shape of the marginal cell; (4) the nearly bare posterior half of the wings; (5) the structure of the nugnes, which is essentially the same as in *Culex*, but the larger claw on the anterior tarsi of the male always has two teeth; (6) the long antennae of the \circ , usually longer than the probose is.

Ludlowia was founded on Mimomyia (?) chamberlaini, Ludlow, and Radioculex on R. clavipalpus, Theo. An examination of the types, however, shows that these are in reality one and the same species. Hispidimyia hispida, Theo., is very little different from L. chamberlaini, and Megaculex albitarsis, Theo., has all the essential characters of the genus.*

Table of species of Ludlowia.

1.	Head with mostly narrow-enrved and upright-forked scales in middle: last two joints of hind tarsi white (subgenus Megaculer)	2.
	Head mostly covered with flat scales, except for a very small area	
	behind; last two joints of hind tarsi not white (subgenus	
	Ludlowia	3.
2.	Large species, 7-8 mm. in length plum	osa.
	Small species, 3 mm. in length pincer	rna.
3.	Legs narrowly banded, last hind tarsal joint whitish (chamberlaid	ui)†
	" uniformly dark hisp	ida.

⁶ Since the above was in type I have had an opportunity of examining a male of *Boycia* mimomylaformis, Newst. (vide p. 266), from Derri, N. Nigeria (J. J. Simpson). It evidently belongs to the subgenus *Megaculex* of *Ludlowia*.

[†] This species does not occur in Africa.

1. L. plumosa, Theo., (Culex) Mon. Cnl. I, p. 373 (1901).

Megaculex albitarsis, Theo., Mon. Cnl. 11, p. 25 (1901).

The hind tarsi of the type of L. plumosa are missing, and as Theobold puts a query against his statement that the hind ungues are simple, I assume that they were absent when the insect was described. The only difference I can see between the types of L. plumosa and M. albitarsis is that in the former the unright forked scales of the head are dark, whereas in the latter they are yellow. There is variation in this respect, however, in the small series (four specimens) of M. albitarsis. Theobald incorrectly describes and figures the fore and mid ungues of the \mathcal{O} M. albitarsis as unidentate : the two teeth on the larger claw are quite evident in the type.

S. Nigeria ; Congo Free State ; Uganda : Mashonaland.

2. L. pincerna, Graham, (*Megaculex*) Am. Mag. Nat. Hist. (8) V, 1910, p. 267. Graham makes the same mistake as Theobald in describing the \mathcal{S} ungues. The basal tooth on the larger claw is much more difficult to see than the median one, and is probably overlooked in very many cases. As, however, its presence or absence is not usually a generic character, the oversight (which is very excusable) is not of serious import.

S. Nigeria.

3. L. hispida, (*Hispidimyia*) Theo., Mon. Cul. V. p. 245 (1910).

Specimens have been received at the Museum from Kampala Swamp and Kibanga, Uganda (*Capt. A. D. Fraser*).

Sudan ; Uganda.

Genus BANKSINELLA, Theo.

Mon. Cul. IV, p. 468 (1907).

1.	Costa	spotted wit	h black	and y	ellow;	hind	femora	yellow	with l	olack
		apical rin	g					ри	nctocost	talis.
	33	uniformly d	lark; hi	nd femo	ora all d	lark al	ove			-2.
2.	Probos	seis unbande	ed					1	uteolate.	ralis.
		banded	•••		• • •			te	aeniaro	stris.

1. B. punctocostalis, Theo., Mon. Cul. V, p. 407 (1910). Ashauti.

2. B. luteolateralis, Theo., Mon. Cul. II, p. 71 (1901).

Taeniorhynchus lineatopennis, Ludlow, Canad. Ent. XXXVII, p. 133 (1905).

Banksinella chrysothorax, Theo., (3) Mon. Cul. V, p. 404 (1910). Generally distributed over Africa; Philippine Is.; Perak, Malay States.

- 3. B. taeniarostris, Theo., Mon. Cul. II, p. 405 (1910).
- The hind ungues of this species and of *B. punctocostalis* are simple, as in *B. luteolateralis*, of which this is perhaps a well-marked variety.

Ashanti.

Genus MUCIDUS, Theo. Mon. Cul. I, p. 268 (1901).

1. M. scatophagoides, Theo., Mon. Cul. I, p. 277 (1901).

M. sudanensis, Theo., Third Rep. Welle. Lab. p. 252 (1908).

Tibiae with pale bands at the base, middle and apex, all of about equal breadth. All tarsi with pale basal bands, most marked on the hind pair. Wing-fringe with eight pale spots.

The probose of *M. sudanensis* is said by Theobald to have a white band at the tip ; this is an error, and the statement was evidently intended to apply to the *palpi*, the last joint of which is white. The species is closely related to the Australian *M. alternaus*, Westw., which differs in having the tibiae dark at the apex, with two narrow white bands.

Sudan; Gold Coast; India.

2. M. mucidus, Karseh, Ent. Naehr. 1887, p. 25.

M. africanus, Theo., Mon. Cul. I, p. 274 (1901).

M. grahami, Theo., Mon. Cul. V, p. 127 (1910).

Fore and mid tibiae narrowly white at base, broadly white at apex. Hind tibiae narrowly white at base and apex and with a very narrow indistinct white band in the middle. Fore and mid tarsi all yellowish. Wing-fringe usually with seven pale spots.

The type of M. grahami is a dark specimen, in which the pale fringe spots are indistinct, only four being clearly visible, and traces of two others. Apart from this it does not differ from M. mucidus. The type of M. africanus has seven pale fringe-spots, not five as stated by Theobald.

M. alternaus does not occur in Africa : the Natal specimen included by Walker in his series of *Culex commoveus* (=M. alternaus) is apparently a variety of *M. mucidus*.

Sierra Leone; Ashanti; S. Nigeria; N. Nigeria; Nyasaland Protectorate; Delagoa Bay.

Genus OCHLEROTATUS, Arrib.

Rev. Mus. La Plata, II, p. 143 (1891).

Culicada, Felt, N.Y. State Mus. Bull. No. 79, p. 391 b (1904).

Culicelsa, Felt, l.c.

Mimeteculex, Theo., Third Rep. Welle. Lab. p. 258 (1908).

Grubhamia, Theo., part.

Coquillett (Science, vol. 23, p. 314, 1906) sinks both *Culicada* and *Culicelsa* under *Ochlerotatus*, and though Theobald (Mon. Cul. IV, p. 14) regards this as a retrograde step, it seems to be inevitable. The present writer is in entire agreement with Coquillett as to the taxonomic value of toothed or simple claws in the female, as all other characters seem to support divisions based on this. A more detailed study may reveal satisfactory characters by which to separate *Culicelsa* and *Ochlerotatus* (*Culicada*), but at present such have not been diseovered. Dyar, using the male genitalia as a basis of classification, doubts whether the two genera can be kept separate. The chief structural difference appears to be that in *Culicelsa* the hind ungues are simple, while in *Culicada* they are toothed; but *O. caliginosus* and *O. ochraceus* have the hind ungues simple in the male, but toothed in the female, so that this character cannot be of very great importance. The genus Ochlerotatus is certainly well founded, and is accepted here in its original sense, with the slight modification necessary for the inclusion of *Culicelsa*. O. albifasciatus, Mcq., the type of the genus, is unrepresented in the British Museum Collection, two specimens so named by Walker certainly being wrongly identified. O. (Leucomyia) scapularis, Rond., is of precisely the same type as the species tabulated below, and is certainly not congeneric with the other species of Leucomyia, which are here included in Culex.

Theobald in publishing *Grabhamia* did not eite a type in any way, and Dyar (Proc. Ent. Soc. Washington, 1905, p. 48) specified *Culex dorsalis*, Mg., as the type; Felt, however, had previously (*loc. cit.*) ehosen *C. jamaiccusis*, Theo., which appears to be a true *Culex*. The genus *Grabhamia*, therefore, is not synonymous with *Ochlerotatus* and *Culicada*, but with *Culex*.

Ochlerotatus is mainly a Palaearetic genus, comparatively few species occurring within the tropies, where, on the other hand, Culex is more largely developed. The Culicelsa group includes (besides C. taeniorhynchus, Wied., the type, and the African species given below), Culex vigilax, Skuse, Taeniorhynchus niger, Giles (which is very close indeed to C. taeniorhynchus, Wied.) and others. C. alboannulatus, Mcq., is a true Ochlerotatus. C. accraeusis, Theo. (neotaeniorhynchus, Theo.) is a Culex. In addition to the differences in ungues and male palpi, Ochlerotatus seems to differ from Culex in the structure of the male antennae, the plumes arising from the dorsal and ventral surfaces of the antenna only, instead of in whorls as in C. pipiens. Whether the two genera can always be distinguished in this way I am unable to say, as time has not allowed the examination of the antennae of very many species.

1.	Hind ungues toothed in both sexes (Ochlerotatus	, s. str.)		2.
	,, ,, simple in ♂		• • •		6.
2.	Tarsi unbanded				3.
	Tarsal joints banded at both base and apex	• • •			5.
3,	Abdomen unbanded, but in Q with very small	media	n basal	pale	
	spots	• • •	1.	cumm	insi.
	Abdominal segments basally banded				4.
4.	Thorax more or less ornamented with pale li	nes ; w	ings w	ithout	
	light scales			2. dent	atus.
	Thorax uniformly sealed ; wings with light	and	dark s	cales	
	(N. Africa and Europe only)		3.	nemor	ดรมร
5.	Banding of tarsi rather broad : abdominal seen	nents w	ith nai	rs of	
	dark patches		in par	4. dars	alis
	Banding of tarsi parrow : abdomen almost all n	ale	5 Jong	ne auro	mene
6.	Hind augues simple in both sexes (<i>Culicelsa</i>)		0. 10119	asyaam	7
0.	toothed in female			• • •	10
7	Legs handed	• • •	• • •	• • •	۰ <i>ـــ</i> ۲ و
**	unbandad	• • •	• • •	• • •	0.
8	Targel joints with anical and bagal banding	• • •	···	···	11.
0,	handling only		0. /	<i>muerna</i>	TSIS.
0	Winga mottled with block and making only	· · ·	• • •	• • •	9.
9.	wings motified with black and white scales; da	rk spec	ies	•••	10,
	,, not mottled; more prownish species; co	stal Iri	nge wn	itish	
10	beyond middle	•••		. hrrsi	ntus.
10,	A buominal segments with apical lateral yellowis	h spots	8. a	urbane	nsis.
	", without such spots		9.	nigerie	nsis.

11. Abdominal segments with complete basal bands ... 10. quasiunivittatus.

tibiae; sentellum with golden-yellow seales ... 14. leucarthrius.

1. 0. cumminsi, Theo. (Culex), Mon. Cul. III, p. 214 (1903).

Culicada mediopunctata, Theo., Mon. Cul. V, p. 304 (1910).

fuscopalpalis, Theo., Mon. Cul. V, p. 307 (1910).

The type of *C. mediopunctata* is only a variety of *O. cumminsi* with mnshally large abdominal spots, traces of which can almost always be seen. The male genitalia of *C. fuscopalpalis* do not differ from those of *O. cumminsi*.

Nyassaland Protectorate ; Uganda ; Ashanti.

2. 0. dentatus, Theo. (Culex), First Rep. Welle. Lab., p. 75 (1905).

Evidently closely allied to *O. cumminsi*, but differs in the banded abdomen. The male is at present unknown.

Abyssinia : Transvaal.

3. 0. nemorosus, Mg., Syst. Besehr. I, p. 4 (1818).

(?) Grabhamia maculosa, Theo., Ann Mns. Nat. Hung. III, p. 105 (1905). Culicada nemorosa, Theo., Mon. Cul. IV, p. 370 (1907).

Several specimens from Algeria (*Rev. E. A. Eaton*) in the British Museum seem to belong to this species. Some are very much paler than others, one having the thorax and abdomen almost entirely buff-coloured, yet I feel sure it is only a variety. This specimen agrees very well with Theobald's description of *G. maculosa*, which makes it seem possible that the latter is only an extreme form of *O. neurorosus*. The corresponding pale variety of *Stegomyia fasciata* (described by Mr. G. A. K. Marshall in the last number of this Bulletin) is common in Algeria, as is an ochroous form of *C. pipieus*.

Algeria.

4. **O. dorsalis**, Mg., Syst. Beschr. IV, p. 242 (1818).

Grabhamia dorsalis, Theo., Mon. Cul. III, p. 251 (1903).

Grabhamia subtilis, Ed. & Et. Serg., Bul. Mus. Paris XI, p. 240 (1905).

willcocksii, Theo., Mon. Cul. IV, p. 294 (1907).

The forms described as *G. subtilis* and *G. willcocksii* only differ in having the pale markings of the abdomen rather more extended than in typical *C. dorsalis*, and are evidently merely pale varieties of Meigen's species.

The ungual formula in all the specimens 1 have seen (including British specimens determined by Mr. Theobald as G. dorsulis and as G. pulcripalpis, and African specimens of G. subtilis and G. willcocksii) is : 32.1-1.1-1.1, Q1.1-1.1-1.1, and it is most probable that errors of observation were made in compiling the table on p. 285 of vol. IV of the Monograph of the Culicidae.

I am not quite certain whether the species at present under consideration is really *C. dorsalis* or *C. pulcripalpis*, Rond., or whether these two names are synonymous. Whichever may be the case, it is certainly the same as our British species, which has always been known as *C. dorsalis*. It is envious, however, that Austen (Entomologist, 1895, p. 228) overlooked the fact that the last joint of the hind tarsi is entirely white or whitish, while Theobald gives this as a character of *C. puleripalpis* (introducing *C. puleripalpis* as British on some specimens which showed the character markedly) but not of *C. dorsalis*. As, however, every specimen in the British Museum, British or otherwise, whether determined as *C. puleripalpis* or as *C. dorsalis*, has the last joint of the hind tarsi whitish, I feel fairly sure that the fact has simply been overlooked. There is in Britain, so far as the National Collection is concerned, only one species of the *C. dorsalis* group, and it seems à priori more probable that this would be Meigen's species than Rondani's.

Mansonia arabica, Giles (J. Trop. Med. 1906, p. 130) is related to O. dorsalis, but can readily be distinguished by the much broader wing-scales, and there are other differences which show it to be distinct. *Calex arabicus*, Becker (Denksehr. k. Akad. Wiss. Math. Nat. kl. LXIX, 2, p. 140, 1910) is probably a homonym, as both appear to belong to the group *Grabhamia*.

Europe : N. Africa.

5. **O. longisquamosus,** Theo. (*Grabhamia*), Ann. Mns. Nat. Hung. III, p. 102 (1905).

The \mathcal{S} and the hind ungues of the \mathcal{Q} of this species have not been described, but it seems to belong to the *O. dorsalis* group, judging from the description. Tunis.

 O. pulcritarsis, Rond., Bull. Soe. Ent. Ital. IV, p. 31 (1872). Calex lencacanthas, Lw., Beschr. Enr. Dipt. III, p. 1 (1873). Calex mariae, Serg., Thèse de Paris, p. 64 (1903). Grabhamia pulcritarsis, Theo., Mon. Cul. III, p. 244 (1903).

The specimens of *C. mariae* in the British Museum agree perfectly with the descriptions of *C. pulcritarsis* (by Fiealbi) and *C. leucacanthus*. The ungual formula in these specimens is $\Im 2 \cdot 1 - 2 \cdot 1 - 0 \cdot 0$, $\Im 1 \cdot 1 - 1 \cdot 1 - 0 \cdot 0$, and not as given by Theobald in Mon. Cul. IV., p. 285; the above is the formula given by Fiealbi for *O. pulcritarsis*. The species is quite distinct from *C. dorsalis*.

Europe ; Algeria.



Fig. 1.—Ochlerotatus hirsutus, Theo., 3; side view of head, to show form and position of palpi.

 O. hirsutus, Theo. (*Culex*), Mon. Cul. I, p. 392 (1901). (Fig. 1.) *Culex transvaalensis*, Theo., Mon. Cul. III, p. 165 (1903).

The probose is has a broad though ill-defined band. This is quite a typical member of the *Culicelsa* group, taking *C. taeniorhynchus*, Wied, (*nec* Arrib.), as the type.

S. Rhodesia : Transvaal.

8. **O. durbanensis**, Theo., Mon. Cul. III, p. 246 (1903). Grabhamia durbanensis, Theo., *l.c.*

ocellata, Theo., Mon. Cul. V, p. 284 (1910).

The type of G. durbanensis is a small specimen, but evidently the same as G. ocellata.

Natal; Delagoa Bay.

9. O. nigeriensis, Theo. (Grabhamia), Mon. Cul. V, p. 281 (1910).

Rather closely resembles *O. rigilax*, Skuse, from Australia and *O. nocturnus*, Theo., from Fiji, but has much more numerous light scales on the wings. The σ of *C. vigilax* has only a protuberance representing the tooth on the

larger claw of the mid tarsi ; the J of C. nigeriensis is not yet known.

S. Nigeria ; N. Nigeria ; Uganda ; Nyasaland.

 O. quasiunivittatus, Theo. (Culex), Mon. Cul. II, p. 32 (1901). Culiciomyia dalzieli, Theo., Mon. Cul. V, p. 234 (1910).

An obscure species, and apparently rare, as only three specimens have been received at the British Museum which can be referred with certainty to it : a male and female from Salisbury, Mashonaland, and a female from Katagum. N. Nigeria. It is not a *Culiciomyia*, and though differing considerably in appearance from the three following species, must be included here on structural characters. It could hardly be mistaken for *C. univittatus*, with which Mr. Theobald has compared it.

N. Nigeria; S. Rhodesia; Natal.

11. **O. lateralis**, Mg., Syst. Beschr. I, p. 5 (1818).

Culicada lateralis, Theo., Mon. Cul. V, p. 310 (1910).

This European species has been recorded also from Algeria. It is very distinct and unlike any other European or African species with the exception of *O. ornatus*, Mg.

12. O. caliginosus, Graham (*Culex*), Ann. Mag. Nat. Hist. (8) v, p. 268 (1910). Resembles *O. quasinnivittatus*, but the hind ingues of the \Im are toothed and the head-scaling is different.

S. Nigeria.

13. **O. ochraceus**, Theo. (*Culex*), Mon. Cul. II, p. 103 (1901).

Culex pallidostriatus, Theo., Mon. Cul. IV, p. 410 (1907).

" parascelos, Theo., Rec. Ind. Mus. IV, p. 18 (1910).

Mimeteculex kingii, Theo., Third Rep. Welle. Lab. p. 258 (1908).

This species can be very easily recognised by the general yellowish eoloration, legs striped with black and yellow, and the unequal hind ungues in the \mathcal{J} . Similar leg-markings are however to be found in *Culex theileri* (= pettigrewii); whilst in *Pecomyia* and *Reedomyia lowisii* the $\mathcal{J}\mathcal{J}$ have unequal hind ungues. I can see no difference between African and Indian specimens; the species, however, appears to be more common in Africa than in India. The unequal hind ungues of the male do not seem to the writer to form a sufficient reason for excluding the species from this genus.

N. Nigeria ; Sudan ; S. Rhodesia ; India ; Ceylon.

14. 0. leucarthrius, Speiser (Culex), Kilimandjaro-Meru Exp., 1905-6, Dipt. Orth., p. 43 (1909).

Dr. Sjöstedt very kindly lent me the type of this species, and I can add the following notes to Dr. Speiser's description :

Head, bearing golden-yellow narrow-enrved seales in the middle.

Thorax black, with brown seales, a short line of golden seales on the lateral margin in front, and a similar short line on each side of the middle line in front of the scutellum. Sentellum covered with narrow-curved bright golden-yellow scales : bristles golden.

Abdomen blackish, the segments with small lateral pale basal spots.

The hind ungues are missing, but it seems to be an Ochlevotatus; it is quite distinct from anything I have seen. The golden-scaled sentellum and the distinct pale band on the apex of the hind tibiae distinguish it at once,

German East Africa.

Genns THEOBALDIA, Neven-Lemaire.

C. R. Soc. Biol., 1902, p. 1331.

Culiseta, Felt, New York State Mus. Bull. 79, p. 391e (1904).

This seems to be a well-marked genus. The claws and antennae are as in Culex, but the male palpi have the last two joints swollen, the terminal porrect and longer than the penultimate, being thus somewhat intermediate in structure between Culex and Ochlerotatus. The genus includes, besides the species mentioned in Theobald's monograph, Culiseta absobrina, Felt, and Culicada morsitans, Theo. The spotting of the wings, as might be expected, is not a constant generic character. Two species have been recorded from Africa.

Thorax with narrow white lines spathipalpis. without white lines annulata, 1. T. spathipalpis, Rond., Bull. Soc. Ent. Ital. IV, p. 31 (1872).

Very variable in size, but easily recognised by the white lines on the thorax and the banded and spotted legs, which have a whitish ring just before the tip of The spots on the legs recall *Culex tigripes*, but the two can hardly be the femora. confused.

Cape Colony; Natal: Transvaal; N. Africa: Madeira; Europe.

2. T. annulata, Schrank, Beit. z. Naturg. 97, 70 (1776).

Algeria.

Genus TAENIORHYNCHUS, Arrib.

Rev. Mns. La Plata, II, p. 147 (1891).

Mausonia, Blanehard, C.R. Soc. Biol. 53, p. 1046 (1901).

This genus is accepted in its original signification, the principal character separating it from other allied genera being the peculiar formation of the ingues (3 2.0-2.0-0.0 or 1.0-1.0-0.0, Q 0.0-0.0-0.0). The wing-scales vary greatly in width from the very broad ones in the type T. titillans, Wlk. (=taeniorhynchus, Arrib., nec Wied.) to the narrow ones of T. aurites and its allies. The author fully agrees with Messrs. Dyar and Knab (Ent. News, July, 1910) that the genus Mansoula was a composite one, the Old World species belonging to a distinct genns, while T. titillans, T. fasciolatus, etc., are congeneric with the species erroneously added by Theobald to Goeldi's genus Chrysoconops; it also

seems probable that Dyar's Coquillettidia is not properly separable from Taeniorhynchus. The type species of the genus Chrysoconops, C. fidous, Wied., has no connection with the African and Oriental species associated with it since the genus was founded; it does indeed resemble them in colour, but in structure it is nearer to Ochlerotatus than to Taeniorhynchus. In the present writer's opinion Culex titillaus, Wlk., must be taken as the type of Arribalzaga's genus, because it is quite evidently the species which he describes, although he wrongly determined it as C. taeniorhynchus, Wied. On this view, which is the reverse of that held by Dyar and Knab, Mansonia becomes a synonym of Taeniorhynchus since C. titillans was also the type of Mansonia.

If C. taeniorhynchus, Wied., be taken as the type, Arribalzaga's eareful and complete description of *Taeniorhynchus* becomes inaecurate in many important particulars; this course, if adopted, would necessitate sinking *Culicelsa*, Felt, as a synonym of *Taeniorhynchus*, and produce the anomaly of a genus none of whose members possessed the generic characters.

1. Violet-black species, thorax with pale scales in front	metallicus.
Abdomen and legs yellow or partly yellow	2.
2. Tibiae all black, abdomen all yellow	nigrithorax.
,, yellow, black-tipped, hind pair with a black ring in r	niddle <mark>3.</mark>
3. Wing seales both dark and light	4.
,, ,, all golden yellow	3. aurites.
4. Thorax with the anterior two-thirds rather densely elothed	with
pale golden seales; joints of hind tarsi black at tips only	annettii.
Thorax uniformly clothed with sparse golden scales ; joints o	f hind
tarsi with broad black apieal bands	5.
5. Thoracic integriment black, abdomen all yellow	cristatus.
", ", brown, abdominal segments with apieal	dark
,, ,, brown, abdominal segments with apieal bands ,, ,, ,, ,, ,, f	dark <i>uscopennutus</i> ,

? Culex nigrochaetae, Theo., J, Mon. Cul. II, p. 60 (1901).

Banksinella metallicus, Theo., Mon. Cul. V, p. 408 (1910).

Taeniorhynchus violaceus, Theo., Third Rep. Welle. Lab., p. 262 (1908).

Chrysoconops nigra, Theo., Mon. Cnl. V, p. 434 (1910).

A very distinct species, easily recognised by the sub-metallic violet colour of the abdomen, femora and tibiae, and the shining whitish area on the front of the thorax.

Sudan ; S. Nigeria ; Angola ; Uganda.

- 2. T. nigrithorax, Theo. (Chrysoconops), Mon. Cul. V, p. 439 (1910). Angola.
- 3. T. aurites, Theo., Mon. Cul. II, p. 209 (1901).

Chrysoconops anarites. Theo., Mon. Cul. IV, p. 493 (1907).

Somewhat variable, some specimens showing dark apical patches on some of the abdominal segments, others having the abdomen entirely yellow. The anterior legs are sometimes without any dark scales, sometimes with the tips of the joints black and the whole of the last two tarsal joints brownish. T, availables are sometimes without any dark scales brownish.

1.

has been wrongly recorded from the Malay States : the specimens are T, ochraceus, which differs from T, aurites in having no median black ring on the hind tibiae.

Sudan ; Uganda : S. Nigeria.

4. T. annettii, Theo., Mon. Cul. II, p. 205 (1901). Chrysoconops annettii, Theo., Mon. Cul. IV., p. 491 (1907).

Chrysoconops pseudoconopas, Theo., Mon. Cul. V., p. 443 (1910).

The specimens of C, pseudoconopas have the thorax very much rubbed, but I can see no difference between them and the types of C, annettii. All the specimens I have seen of C, annettii have the distinguishing characters given by Theobald for C, pseudoconopas. The dark scales on the wings occur chiefly in two tranverse bands, one before the tip of the wing and including the bases of the fork-cells, the other nearer the base. These bands are more conspicuous in some specimens than in others.

Uganda ; S. Nigeria.

 T. cristatus, Theo., First Rep. Welle. Lab., p. 78 (1904). Chrysoconops cristatus, Theo., Mon. Cul. IV., p. 491 (1907). Sudan; N. Nigeria; S. Nigeria; Congo Free State (Katanga).

 T. fuscopennatus, Theo., Mon. Cul. III, p. 265 (1903). Chrysoconops fuscopenatus, Theo., Mon. Cul. IV., p. 492 (1907). Culex drymoecius, Speiser, Kilimandjaro-Mern Exp., Dipt. Orth. p. 42 (1909).

Dr. Sjöstedt very kindly lent me a specimen of *C. drymoecius*, and it proved to be quite a typical *T. fuscopennatus*, though it certainly did not show the black abdominal spots mentioned by Dr. Speiser in his description. As, however, one or two males in the British Muscum collection show these spots rather indistinctly, I think *C. drymoecius* can only be a variety of *T. fuscopennatus*.

Uganda.

Genus MANSONIOIDES, Theo.

Mon. Cul. IV., p. 498 (1907).

Mansonia, Blanchard, C. R. Soc. Biol., LIII, p. 1046 (1901) (part).

As suggested by Messrs. Dyar and Knab, and noted above, the Old-World species of "Mansonia" cannot be kept in Taeniorhynchus. They resemble that genus in the peculiar structure of the elaws, but the male palpi are so different that they are obviously distinct. The penultimate joint is thin and curved upwards, as in Culex, but the terminal joint is very short— $\frac{1}{8}$ to $\frac{1}{4}$ as long as the penultimate, and bent downwards at an angle with it. The terminal joint, in all the males I have seen (M. uniformis, M. annulipes and M. annulifera) is entirely covered with white scales. The resemblance in all structural details, and in many minor characters as well, between M. annulifera and M. uniformis is so great that I do not think it is possible, in this case at least, to regard the presence of flat or narrow-curved scales on the scutellum as a generic character. Probably Etorleptiomyia is a synonym, but as the male of E. mediolineata is unknown, I have adopted the later name Mansonioides for this genus.

1.	Tarsi basally banded			2.
	First three tarsal joints apically banded			mediolineata.
2.	Abdomen with narrow apieal pale bands,	sente	ellum	
	flat-sealed (?)			nigra.
	Abdomen with apical lateral pale spots			uniforms.
2136	7			E 2

- M. (?) mediolineata, Theo., First Rep. Welle. Lab., p. 71 (1904). Etorleptiomyia mediolineata, Theo., I.c. Sudan; Transvaal.
- M. (?) nigra, Theo., Second Rep. Welle. Lab., p. 80 (1906). Mansonia (?) nigra, Theo., *l.e.*

It is not easy to see why Theobald did not include this species in *Etorleptiomyia*, as the heart-shaped scales are even more prononneed than in *M. mediolineata*. The unique type is now very badly damaged, but is clearly distinct from the other two species here associated with it.

Sudan.

3. M. uniformis, Theo. (*Panoplites*), Mon. Cul. II, p. 180 (1901) (fig. 2). The following are included here :--

Mansonia africana, Theo., Mon. Cul. II, p. 187 (1901).

- " major, Theo., Mon. Cul. III, p. 270 (1903).
- " uigervina, Theo., Mon. Cul. V, p. 450 (1910).

In the form *africana* the two greyish stripes which are present on the thorax of *M. uniformis* (typical form) are more or less broken up into spots. The names *major* and *nigerrima* have been applied to forms in which the reddish-brown



FIG. 2.—Mansonioides uniformis, Theo., & ; side view of head. Note structure of palpi.

colour of the thorax is replaced by blackish, and the pale bands of the legs are narrower than usual. These forms, however, grade into one another, and it seems best to regard them as only varieties of M. uniformis.

Common throughout the Ethiopian, Oriental, and Australian regions.

Genus CULICIOMYIA, Theo.

Mon. Cul. IV, p. 227 (1907).

Neomelanoconion, Theo., Mon. Cul. IV, p. 514 (1907) (male only). Pectinopalpus, Theo., Ann. Mag. Nat. Hist. (8) V, p. 375 (1910).

C. nebulosa, Theo., Rept. Liverp. S. Trop. Med., Mem. IV, App., p. x. (1901).
 (?) Culex invenustus, Theo., op cit., p. ix. (1901).

Culex cinereus, Theo., Mon. Cul. II, p. 58 (1901).

- " nigrochaetae, Q, Theo., Mon. Cul. II, p. 60 (1901).
- , pseudocinercus, Theo., Mon. Cul. II, p. 62 (1901).
- " freetownensis, Theo., Mon. Cul. II, p. 69 (1901).

Neomelanoconion rima, &, Theo., Mon. Cul. IV, p. 514 (1907).

Pectinopalpus fuscus, Theo., Ann. Mag. Nat. Hist. (8) V, p. 375 (1910).

Culiciomyia uniformis, Theo., Mon. Cul. V, p. 230 (1910).

Dr. W. M. Graham first called the writer's attention to the probability of Pectinopalpus fuseus and Culiciomyia freetomnensis being identical, and has suggested this in the "Bulletin of Entomological Research" (July, 1911, p. 132). At his suggestion I made preparations of the genitalia of a number of males, choosing specimens as different in size and in the amount of white on the abdomen as I could find. In none of these preparations could I see any specific differences; the smaller specimens were much less strongly chitinised, but this is only what might be expected in starved individuals. I also examined some hundreds of pinned adults, and though these varied greatly in size and in the extent of the whitish, lateral, abdominal spots, every gradation could be found between the varions forms. In one case about 150 very small specimens were bred from the water in a pie-dish by Dr. T. F. G. Mayer, in Southern Nigeria, their size evidently being due to semi-starvation in the larval period. I have therefore come to the conclusion that there is only one variable African species of Culiciomyia (C. dalzieli, Theo., is a synonym of Ochlerotatus quasinnivittatus, Theo.), the forms or varieties of which are :--

- a. nebulosa. Medium-sized specimens with obscure abdominal spots.
- β . cinerea. Large specimens with distinct pale apieal lateral spots on the abdominal segments.
- γ . pseudocinerea (= uniformis, ? = invenusta). Large specimens with no pale abdominal spots.
- 8. freetownensis. Small specimens with abdominal spots.
- E. fusca. Small specimens without abdominal spots.

As it would frequently be impossible to say whether a specimen were large or small, or even whether abdominal spots were present or not, since these are so variable in size, it does not seem advisable to retain these names in general use. There is some variation in the length of the fork-cells : two specimens from Prince's Island, Gulf of Guinea (Dr. W. G. Ansorge) have the first slightly shorter than the second. This is one of the many species which not infrequently have a bright green colour on the thorax or legs : this variation may possibly be due to something in the food of the larva.

In the Monograph of the Culicidæ (V, p. 239), referring to C. freetownensis, Theobald says "Perfect specimens show the venter of the abdomen to be white, except the apical segment, which is deep black." This remark only applies to some specimens wrongly determined as C. freetownensis, which are really Eumetanomyia inconspicuosa; the venter of C. nebulosa is all grey. E. inconspicuosa Q is very much like C. nebulosa to the naked eye, but has a black, shiny thorax, and a flat-scaled head. The thoracic integrument of C. nebulosa and also the scales of the thorax, have a characteristic dull greybrown appearance. The distinctions between C. nebulosa Q and Protomelanoconion fuscum Q have already been pointed ont.

The type of *C. nigrochaetae* is not now in the British Mnsemn, but it seems, from the description of the Q, to have been only *C. nebulosa*. The number of

unid-sentellar bristles, as usual, is variable in this species, and as a matter of fact there are only 6, not 8, in the type of *C. cinerea*. I cannot help thinking, however, that the \mathcal{J} described as *C. nigrochaetae* was in reality *Taeniorhynchus* metallicus. Calex invenustus is probably the same species, but as I have not seen the type I have used the name nebulosa, which was published a page later in the same paper as *C. invenustus*. *C. invenustus* is said to have the thorax covered with flat scales (a character which is not found in any *Culex* I have seen), and to have the fore and mid femora swollen. If the former statement is an error, and the latter simply due to the femora having become flattened (a frequent occurrence), there cannot be much donbt that the specimen described as *C. invenustus* was a *C. nebulosa* var. pseudocinerea.

C. nebulosa appears to be one of the commonest of the CULICIDAE in West Africa, and extends into the Congo Free State, Uganda and Mashoualand.

Genns CULEX, Linn.

Syst. Nat. Ed. X (1758).

Culex, as now restricted, is one of the easiest genera to recognise in the male, but the females are sometimes not easily distinguished from those of Taeniorhynchus, etc. The genus includes only those species whose males have palpi and ungues similar in structure to those of C. pipiens. The mere upward curvature of the palpi may not be considered of importance, but as far as the writer has seen it seems to be perfectly constant, and when once understood forms the easiest character by which to distinguish a species of true *Culex* from one belonging to an allied genus. The structure and form of the \mathcal{J} palpi will be nuclerstood by a glance at the figure of C. quasigelidus \mathcal{J} (p. 258). The genus Culex, in this sense, includes Lutzia, Theo., Aporoculey, Theo., Leucomyia, Theo. (except L. scapularis), Melanoconion, Theo., Heptaphlebomyia, Theo., Pseudoheptaphlebomyia, Ventr., and some species of Tacniorhynchus, Theo. (T. tenax and T. ager), of Grabhamia (G. jamaicensis, etc), of Culicada (C. quasimodesta and C. bicolor) and of Culicelsa (C. accraensis and C. annulirostris) besides some small genera, such as Lasiocomps and Ocaleomyia, which were only founded on misconceptions : as regards the last named, contiguous eyes are found in nearly all the CULICALES, and I believe in most of the METANOTOTRICHA with the exception of *Eretmapodites*.

1.	Proboscis and tarsi with pale bands, those on tarsi including both	
	sides of joints (fig. 4, a)	2.
	Proboseis and tarsi unbanded	7.
$\dot{2}.$	Thorax with pale scales on the anterior two-thirds, or at least	
	with a transverse pale band behind the middle (Leucomyia, Theo.)	
	(fig. 5, a)	3.
	Thorax with almost uniform scales, at most with a pair of pale	
	spots, abdominal segments with complete basal white bands	5.
3,	Femora and tibiac with rows of sharply defined whitish spots	
	(fig. 4, a) 1. quasigelid	us.
	Femora and tibiae not spotted	4.
4.	Abdomen ahnost unicolorous dark brown 2. consimi	lis.
	Adominal segments with both median basal and lateral apical	
	triangular pale spots 3. annulion	ris.

5.	Middle tibiae with a whitish lateral stripe (fig. 4, b); abdominal bands broadest in middle ; hand of probases broad and ill-defined
	Janus Stoatest II militate; Saint of probosits Stoat and In-demice
	Middle tibine unstrined : abdominal bands of equal breadth
	throughout - hand of probosels parrower and more clearly
	defined
G	Fourse not marhled 5 thalassing
0.	morbled (i.e. with dark and light scales interported)
	, marmed (i.e., with dark and light scales interspersed)
7	I area spacing (wearly 7 mm, or more in length) : low swetted
1.	Targe species (usually 7 min. or more in length); legs spotted
	Malium sized a small sussing laws not sputted
0	The string of small species; legs not spotted o.
o.	Legs striped somewhat as in <i>Ochterolatus ochraceus</i> (on the fore
	legs the paie stripe is more or less broken up into spots, recalling
	Cutex trigripes) (fig. 4, c) 8. thetaevi.
0	Legs not so marked \dots \dots \dots \dots \dots \dots \dots \dots y .
9.	Tible striped with whitish (fig. 4, a) much as in <i>Callex automi</i> ;
	seventh vem more distinct than usual, and generally bearing a
	few scales (Heptaphlebomyna, Theo.) 9. univitatins.
	Legs uniformly brownish (except for pale knee-spots) 10.
10.	Abdominal segments with pale markings basally 11.
	,, ,, ,, apically 19.
11.	Abdominal segments with basal banding in both sexes 12.
	", ", with basal lateral pale spots, in the \mathcal{J} often united
	in the middle line to form an irregular-shaped band 14.
12.	Larger species, abdominal bands yellowish 10. pipieus group.
	Smaller species, abdominal bands white 13.
13.	Thoracic scaling uniformly reddish-brown 11. deceus.
	", ", partly bronzy-brown and partly brassy, the latter
	often predominating 12. simpsoui
14.	Thorax all pale-scaled, integument with lateral pale areas in front
	(fig. 5, b); abdominal spots large in both sexes 13. pruiua.
	Thorax not all pale-scaled, integument uniformly dark 15.
15.	Thorax with two distinct pale areas near the middle (fig. 5, c)
	14. oruatothoracis.
	, with uniform scaling or nearly so 16.
16.	Venter entirely whitish 15. guiarti.
	,, with dark bands on the apices of the segments 17.
17.	Bases of fork-cells in Q equidistant from wing-base 16. grahami.
	Base of first fork-cell nearer wing-base than that of second 18.
18.	Larger species (usually 5-6 mm.); thorax with reddish brown scales
	17. zombaensis.
	Smaller species (usually 3-4 mm.); thorax with dark brown scales
	18. invidiosus.
19.	Abdomen with complete apical bands on at least some of the
	segments 20.
	Abdomen with apical lateral spots 22.

... ... 22.

F. W. EDWARDS-THE AFRICAN

20. Ter:	minal j	oint of	Q palp	i short :	and thic	k		19. sal	isburie	ensis.
	99		•9	long a	nd thin	• •				21.
21. Abd	loninal	l segmei	nts 2-7	with ap	ical pale	e band	8	20	. pilif	erns.
	• •	••	2-4	only w	ith com	plete	apical	pale		
			b:	ands		· · · ·		· 2	I. insi	gnis.
22. Tho	racic i	ntegunio	ent black	kish. se	ales blac	ekish		•••	22.	r rima.
	94	,,	redd	ish	•••		• • •			22.
23. Scal	es and	bristles	of the	rax red	dish bro	WII		2:	3. serg	enti.
95		9 •	• 7	bla	ck	•••		24.	rnbin	otus.
	Ø									



Fig. 3.—Culex quasigelidus, Theo. 5 ; side view of head. Note the long, hairy, upwardly curved palpi.

- C. quasigelidus, Theo., Mon. Cul. 111, p. 181 (1903) (figs. 3, 4a & 5a). Lasioconops poicilipes, Theo., Rept. Liverp. S. Trop. Mcd., Mcm. IV. App. p. ix (1901); Mon. Cul. 111, p. 236 (1903).
 - Culex taeniorhynchoides, Giles, J. Trop. Med. 1904, p. 369.
 - Psendoheptaphlebomyia madagascariensis, Ventrillon, Bul. Mus. Paris, X1, p. 427 (1905).
 - Tachiorhynchus tenax var. machlipes, Theo., First Rep. Welle. Lab., p. 79 (1905).

Approcules punctipes, Theo., Mon. Cul. IV, p. 316 (1907).

Lencomyia quasigelida, Theo., Mon. Cul. IV, p. 372 (1907).

The only species with which this can be confused is *C. tigripes*, which is larger, and has an unbanded probose is and more numerous spots on the legs: the thoracic scaling is also quite different. *C. quasigelidus* has a very distinct appearance, as might be supposed from the fact that three genera have been erected for its special benefit. One of these, *Lasioconops*, was founded on a specimen which had a number of scales of some Lepidopteron adhering loosely to its abdomen. Some specimens of *Pseudoheptaphlebomyia* sent to the British Museum by Dr. Ventrillon are undoubtedly this species. In this and the two following allied forms the pale front two-thirds of the thorax is often more marked in the \Im than in the Q. Gambia; N. Nigeria; S. Nigeria; Sudan; Uganda: Nyasaland Protectorate; Congo Free State; Angola: Madagascar.

 C. consimilis, Newstead, Ann. Trop. Med. I, p. 23 (1907). Calex tigripes var. consimilis, Newstead, l. c. Culex pseudoananlioris, Theo., Mon. Cul. V. p. 333 (1910). Taeniorhyachus tenax, Theo., Mon. Cul. III, p. 258 (part).

This is very closely allied to *C. quasigelidns*, and may perhaps be only a variety of that species : the difference in the leg marking, however, is so very striking that I prefer to consider them distinct. The four specimens of *C. pseudo-annulioris* in the British Museum collection are all rubbed almost bare (they must have been described in that state), but 1 believe they can be safely included under *C. consimilis*. This species is no doubt the African representative of the Oriental *C. ager*, but it is difficult to accept Mr. Theobald's opinion that the two forms are conspecific : the true *C. ager* (= *Taeniorhyachus tenax*) is not African. Of the Oriental species there are two varieties :—

- a. C. ager, Giles, Entomologist, XXXIV, p. 196 (July 1901). Taemorhynchus ager, Giles, l. c. Taemorhynchus tenar, Theo., Mon. Cul. II, p. 198 (Nov. 1901). Oculeomyja sarawaki, Theo., Mon. Cul. IV, p. 515 (1907).
- β. C. ager var. sinensis, Theo., Mon. Cul. III, p. 180 (1903). Lencomyia gelida var. sinensis, Theo., l. c. Lencomyia sinensis, Theo., Mon. Cul. V, p. 313 (1910). Grabhamia ambiguo, Theo., Mon. Cul. III, p. 248 (1903). Grabhamia taeniarostris, Theo., Mon. Cul. IV, p. 299 (1907).

The var. *sinensis* is distinguished from the type form by the presence of basal as well as apical yellow bands on the abdominal segments.

In Mr. Newstead's series of *C. consimilis* are specimens of the preceding species, but the type has no spots on the legs.

Sierra Leone : Ashanti ; S. Nigeria ; Congo Free State : Sudan : Uganda ; Natal.

3. C. annulioris, Theo., Liverp. S. Trop. Med., Mem. V, App. (1901).

This species is really very distinct, but has been confused with *C. duttoni*, one of the types (\mathcal{S}) of *C. hirsutipalpis* (= *duttani*) actually being a specimen of *C. annulioris*. The markings of the thorax and abdomen, and the absence of the stripe on the tibiæ, will suffice to distinguish the present species from *C. duttani*. Gambia; S. Nigeria; Angola; S. Rhodesia; Transvaal.

- 4. C. duttoni, Theo., Rept. Liverp. S. Trop. Med. Mem. IV, App. p. v. (1901). C. dissimilis, Theo., Mon. Cul. I, p. 376 (1901).
 - C. hirsutipalpis, Theo., Mon. Cul. I, p. 378 (1901).
 - C. anarmostus, Theo., Rept. Liverp. S. Trop. Med., Mem. X, App. p. vi. (1902); Mon. Cul. III, p. 170 (1903).
 - (?) C. condyladesmus, Grünb., Zool. Anz. 29, p. 385 (1905).
 - C. allovirgatos, Graham, Ann. Mag. Nat. Hist. (8) V, p. 264 (1910).

The pale stripe on the tibiæ of this species is sometimes rather obscure, but in conjunction with the broad band on the proboseis, the two pale spots on the thorax (not always visible), and the rounded abdominal bands, it makes the species an easy one to identify. *C. univittatus* has the same spotted thorax, striped tibia and rounded abdominal bands, but differs in the unbanded probose and legs. The tibial stripe, too, in the last-named species is most distinct on the hind legs, while in *C. duttoni* it is particularly conspicuous on the middle legs. In the Q type of *C. dissimilis* I cannot make out the markings of the tibia, and the fork-cells are different; in fact this specimen looks much more like *C. thalassius* than *C. duttoni*, but since the males associated with it are undonbtedly all *C. duttoni* I have included *C. dissimilis* here as a synonym. I feel all the more confident in doing this, since Theobald (Entomologist, 1908, p. 107) has already pointed out that *C. hirsutipalpis* is only a large form of *C. dissimilis*. *C. hirsutipalpis* differs from the typical form in having some pale scales at the apieces as well as at the bases of the abdominal segments, but there is no doubt



Fig. 4.—Legs of Culex: (a) Hind leg of C. quasigelidus, Theo.; (b) mid leg of C. duttoni, Theo.; (c) mid leg of C. theileri, Theo.; (d) hind leg of C. univittatus, Theo.

that this is only a variation, and it is hardly worth while retaining a separate name for it. *C. anarmostus* was described from a small specimen; this species, like many other mosquitos, varies very greatly in size. One specimen shows a remarkable abnormality in the neuration: the second vein is unbranched, and the upper branch of the fourth vein is only represented by a stump, so that the forkcells are absent. *C. condylodesmus* is probably a synomym, but the description does not quite agree, as the band of the probose of the female is described as being broader and *more distinct* than in *C. dissimilis*.

Common throughout West Africa ; Nyasaland ; Transvaal.

- 5. C. thalassius, Theo., Rept. Liverp. S. Trop. Med., Mem. X, App. p. vii (1902); Mon. Cul. III, p. 168 (1903).
 - Culex bifoliata, Theo., J. Econ. Biol. I, p. 31 (1905).

Calicelsa accraensis, Theo., Mon. Cul. V, p. 317 (1910).

" ucotaeuiorhynchus, Theo., Mon. Cul. V, p. 320 (1910).

There is very little doubt that *C. bifoliata* is the same as *C. thalassins*, which is certainly conspecific with *C. accraensis*. The proboscis in both the specimens of *C. bifoliata*, contrary to Theobald's statement, is distinctly banded. I have examined the σ genitalia in these two specimens, and also in a specimen from Accra, and can only detect one "foliate plate" in any of them. There appears to be no difference in the thoracic marking between *C. accraeusis* and *C. neotaeniorhynehus*—at any rate not sufficient to warrant the retention of the latter name as a distinct variety. This name is rather misleading, as *C. thalassius* does not belong to the *Culicelsa* group, the type of which is *C. taeniorhynchus*, Wied.

Gambia; S. Nigeria; Transvaal; Delagoa Bay.

6. C. somaliensis, Neveu-Lemaire, Arch. Parasit. 10, p. 254 (1906).

C. salus, Theo., Third Rep. Welle. Lab. p. 256 (1909).

C. salsus, Theo., Mon. Cul. V, p. 338 (1910).

Very much like C. thalassius, but the first fork-cell is shorter and the femora are marbled. It bears an extremely close resemblance to the Indian C. microannulatus, and is quite possibly only a form of that species. As, however, I do not feel confident that they are the same—there is some difference in the shape and colour of the thoracic scales—I have not adopted the name microannulatus for the African species. The description of C. somaliensis agrees very well with specimens of C. salns, a large number of which have been received by the Entomological Research Committee from Somaliland.

- 7. C. tigripes, Grandpré, Les Monstiques (1900).
 - (?) C. coucolor, R.-D., Mém. Soc. d'Hist. Nat. Paris IV, p. 405 (1825).
 - C. macalierura, Theo., Mon. Cul. 11, p. 34 (1901).
 - C. tigvipes var. fusca, Theo., Mon. Cul. V, p. 394 (1910).

This species is very distinct from all other Afriean *Culex* on account of its large size and spotted legs. It is, however, closely related to the Oriental *C. concolor*, the only constant difference I can detect being in the relative length of the fork-cells : in *C. tigripes* Q the base of the first fork-cell is nearer the base of the wing than that of the second, while in *C. concolor* the bases of both are almost equidistant from the base of the wing, that of the second fork-cell being if anything nearer the wing-base than that of the first. *C. tigripes* is also on the whole a darker insect, with less yellow on the abdomen, though there is a great deal of variation in this respect in both forms, some specimens of *C. concolor* being as dark as any *C. tigripes*, and having hardly any yellow scales on the abdomen. The last joint of the σ palpi in *C. tigripes* is generally dark, while in *C. concolor* it is nearly always light. Both forms have the same legmarkings, these being subject to some variation. Since there is this constant difference between the two forms (in neuration) I think *C. tigripes* must be regarded as a distinct species. To my mind it is an absolutely typical *Culex*.

It is enrious to notice, that when a species is common to the Ethiopian and and Oriental regions, or when a pair of closely allied species occurs in the two regions respectively, the African form is often much darker in colour than the Eastern. The phenomenon does not seem to be confined to the CULICIDAE, as I have noticed it in the Tipnlid *Conosia irrorata*, Wied.

The specimens mentioned by Theobald as having been taken in Anstralia belong to the *concolor* form. I have made several preparations of the male genitalia of each form, and can detect no difference whatever; the genitalia are of the ordinary *pipiens* and *fatigans* type. Theobald's *C. tigripes* var. *fasca* (Mon. Cul. V, p. 394) the male genitalia of which I have also examined, includes the darkest forms. It is certainly not, as I at first thought possible, a distinct species. *C. tigripes* is widely distributed over Africa.

8. C. theileri, Theo., Mon. Cul. III, p. 187 (1903).

C. pettigrewi, Theo., Rec. Ind. Mus. IV, p. 15 (1910).

A species very easily recognised by means of the striped femora. The co-type of *C. pettigrewi* in the British Museum exactly resembles *C. theileri*.

Transvaal ; Madeira ; India.

 C. univittatus, Theo., Mon. Cul. II, p. 29 (1901). *Heptaphlebomyia simplex*, Theo., Mon. Cul. III, p. 337 (1903).

Heptaphlebomyia montforti, Ventrillon, Arch. Parasit. IX, p. 448 (1905).

A rather obsence species, until the striped tibiae are noticed. Some specimens recall *C. duttoni*, but they can always be distinguished by the unbanded probose is and legs. The British Museum series of *H. simplex* consisted of 3 \bigcirc and 1 \bigcirc *C. univittatus* (including type \bigcirc of *H. simplex*) and one \bigcirc *C. decens* (labelled as type of *H. simplex*, and figured in Mon. Cul. IV, p. 533). The co-types of *H. montforti* from Madagascar, which are in the British Museum collection, are certainly also *C. univittatus*. I do not believe the species is separable generically from *Culex*. The whitish spot on the apex of the tibiae is not more conspicuous in this species than in other members of the genus.

N. Nigeria ; Angola : S. Rhodesia ; Madagascar.

10. C. pipiens, Linn., Syst. Nat. Ed. X, p. 602 (1758).

Included here as varieties are :

i. C. fatigans, Wied., Anss. Zweif. Ins. p. 10 (1828). The stem of the first fork-cell is a little longer than in C. pipiens.

E. Africa; common almost everywhere in other parts of the tropics.

- C. fatigans is said by Grünberg (Zool. Anz. xxix, 1905, p. 390) to be, with Stegomyia fasciata, F., the commonest Culicine in Kamerum and Togo. It seems as though some mistake must have been made, as C. fatigans is unrepresented in any of the large collections of mosquitos which have been received in Eugland from West Africa.*
- ii. C. pallidocephala, Theo., First Rept. Welle. Lab. p. 73 (1904). The scales of the head are paler and the thorax is dark brown instead of reddish brown. The costal margin is darker.

Sudan; British East Africa.

262

^{*} Since the above was written I have seen three specimens from Sokoto, N. Nigeria (Dr. J. M. Dalziel).

- iii. C stochri. Theo., Mon. Cul. IV, p. 419 (1907). The abdominal bands are more rounded, otherwise the specimen is like C. pallidocephala. Nyasaland.
- iv. C. carioannulatus, Theo., Mon. Cul. III, p. 198 (= C. azoriensis, Theo., Mon. Cul. III, p. 210, 1903). This is an almost typical C. pipiens: the females are rather shrivelled, hence the bands on the apical segments of the abdomen are not visible.

Cape Colony : Azores.

v. C. pipiens, L.

N. Africa; Europe; N. America.

11. C. decens, Theo., Rept. Liverp. S. Trop. Med. Mem. IV, App. p. vii, (1901).

C. masculus, Theo., Mon. Cul. II, p. 125 (1901).

C. minutus, Theo., J. Econ. Biol. I, p. 30 (1905).

C. nigrocostalis, Theo., Mon. Cul. V, p. 367 (1910).

C. lividocostalis, Graham, Ann. Mag. Nat. Hist. (8) V, p. 269 (1910).

Heptaphlebomyia simplex ♂, Theo., Mon. Cul. IV, p. 533 (1907).

Theobald states that in *C. masculus* the smaller claw on the fore and mid tarsi of the \mathcal{J} is not toothed; if this is the case (it is difficult to make out from the specimens), it is a very unusual variation. *C. masculus* seems otherwise identical with *C. decens*, which is a small and rather obscure species. Out of 18 specimens in the National Collection only 2 are females.

Sierra Leone ; S. Nigeria ; N. Nigeria ; Sudan ; Uganda ; Transvaal.

12. C. simpsoni, Theo., J. Econ. Biol. I, p. 28 (1905).

Very much like C. decens, differing in the thoracic scaling. Transvaal.

13. C. pruina, Theo., Rept. Liverp. S. Trop. Med. Mem. IV, App. p. viii, (1901).

C. pruinosns, Theo., Mon. Cul. II, p. 332 (1901).

C. pallidothoracis, Theo., Mon. Cnl. V, p. 370 (1910).

Easily distinguished from all other African species by the pale lateral areas on the thoracic integument.

Ashanti ; S. Nigeria.

14. C. ornatothoracis, Theo., Mon. Cul. V, p. 376 (1910).

Very near C. grahami. Both have the female palpi longer than usual; in this species they are quite one-fourth the length of the probosels.

S. Nigeria ; Gold Coast.

15. C. guiarti, Blanchard, Les Moustiques, p. 629 (1905).

C. viridis, Theo. (nec R.-D., 1827), Mon. Cul. III, p. 212 (1903).

C. neuvei, Theo., Second Rept. Wellc. Lab. p. 76 (1906).

The green colour is certainly not a specific character; it not uncommonly occurs in many species, e.g., C. pipiens, C. grahami, C. penina, Culiciomyia nebulosa, and the Tipulid Dicranomyia chorea.

Sudan; Uganda; N. Nigeria; Gold Coast.

16. C. grahami, Theo., Mon. Cul. V, p. 628 (1910).

C. pullatus, Graham (acc Coquillett, 1904), Ann. Mag. Nat. Hist. (8) V, p. 265 (1910).

This and the two following species are extremely closely allied and perhaps should be classed together. Larger series and a closer study are, however, required. There is sometimes in *C. grahami* a more or less distinct rim of white scales to the eyes, recalling that of *Protomelanoconion fuscum*.

Gold Coast ; S. Nigeria ; Uganda.

Uganda; Nyasaland.

17. C. zombaensis, Theo., Mon. Cul. II, p. 143 (1901).

C. quasiguiarti, Theo., Mon. Cul. V, p. 374 (1910).

C. quasigniarti may be distinct, but I can see no character to separate it from C, zounbaensis.



FIG. 5.—Dorsal view of thorax of, (a) Culex quasigelidus, Theo., φ; (b) Culex pruina, Theo., φ, the right half denuded to show the pale anterior patch; (c) Culex ornatothoracis, Theo., φ.

18. C. invidiosus, Theo., Mon., Cul. II, p. 329 (1901).

C. euclastus, Theo., Rept. Liverp. S. Trop. Med., Mem. X. App. viii. (1903).

C. chloroventer, Theo., Mon. Cul. V, p. 373 (1910).

(?) C. aquilus, Graham, Ann. Mag. Nat. Hist. (8) V, 1910, p. 266.

If the males described as *C. aquilus* do not belong to this species, there is no male in the collection.

Gambia : Gold Coast ; Nigeria ; Congo Free State ; Uganda.

C. salisburiensis, Theo., Mon. Cnl. II, p. 113 (1901).
 C. bostocki, Theo., J. Econ. Biol. I, p. 29 (1905).

The type of *C. bostochi* has been lost, but the description leaves practically no doubt that it was only *C. salisburiensis*.

S. Rhodesia ; Transvaal ; Natal.

264

20. C. piliferus, Theo., Mon. Cul. IV, p. 274 (1907). Maillotia pilifera, Theo. l.c.

This species closely resembles a small *Culex*, so is included here, as *Maillotia* only differs from *Culex* in the size of the scales on the head and thorax. \mathcal{J} unknown.

Algeria.

21. C. insignis, Carter, Bull. Ent. Res. II, p. 37 (1911). Culiciomyia insignis, Carter, *l.e.*

Nyasaland ; Uganda ; Congo Free State (Colquilhatville) ; S. Nigeria ; Sierra Leone.

22. C. rima, Theo., Q, Mon. Cul. II, p. 327 (1901).

(Nec Neomelanoconion rima, Theo., J, Mon. Cul. IV, p. 514, 1907).

Melanoconion rimus, Theo., Mon. Cul. III, p. 240 (1903).

Culex rima was founded on female examples, the specimens which Theobald took to be the males are only Culiciomyia nebulosa; the \mathcal{S} of C. rima has not got the peculiar scales on the palpi or the flat scales round the eyes. Since the genus Neomelanoconion was founded on the male specimens in question it is a synonym of Culiciomyia.

Ashanti; S. Nigeria; Congo Free State; Uganda.

23. C. sergenti, Theo., Mon. Cul. III, p. 218 (1903). Algeria.

24. C. rubinotus, Theo., Second Rep. Welle. Lab., p. 76 (1906).

 Λ single specimen, with very few scales left on the thorax or abdomen. Sudan.

African Species not included in the preceding Tables.

C. aegypti, Linn., Hasselquists' Reise nach Palestina, p. 470 (1762). The description is as follows :—" Calex aegypti articulationibus candidis. Magnitudo Culicis vulgaris Linn. Syst. N. 1. Color ex fusco canus. Crura cana cum annulis candidis, parvis, eirea articulationes et in articulis. Puneta candida ad marginem dorsi in corpore sub alis utrinque, plura, longitudinaliter sita. Annulus candidus ad basin thoracis, inter illum et corpus. Linea candida perpendienlaris juxta oculos, utrinque una parva. Locus, Aegyptus, Culice communi rarior."

C. longefurcatus, Beeker, Berl. Mitt. Zool. II, 3, p. 68 (1903). In his description Beeker says "One sees no trace of scaling on the whole body." The name had therefore better be dropped altogether.

C. maculiventris, Meq., Dipt. Exot. Snp. I, p. 7 (1846). This evidently belongs to the *dorsalis* group, but the description is not full enough for purposes of tabulation.

C. rufinus, Bigot (Theo., Mon. Cul. II, p. 169); C. pusillus, Meq. (Theo., Mon. Cul. II, pp. 122, 166); C. pallipes, Meq., nec Mg. (= C. melanovhinus, Giles; Theo., Mon. Cul. II, p. 171). The descriptions of these species are inadequate.

Culicada quasimodesta, Theo., Anu. Mus. Nat. Hung. III, p. 88 (1905), and Culex bicolor, Mg., Syst. Beschr. I, p. 9 (1818), both seem to belong to Culex (s. str.) and apparently to the C. pipiens group. I have not seen specimens.

Culex didieri, C. pygmaeus and C. zeltneri, Neveu-Lemaire, Arch. Parasit. 1906. I have not seen specimens.

C. mundulus, Grünb., Zool. Anz. 29 p. 388 (1905). Said to be closely related to *Culiciomyia nebulosa*, but larger and paler. The description agrees fairly well with the latter species, except that the venter is described as having the segments with pale apical bands.

C. par, Newst. Ann. Trop. Med. 1, p. 25 (1907). From the description this appears to be closely allied to C. consimilis, and is perhaps a rubbed specimen of that species.

C. laurenti, Newst., l.c. p. 24. The abdomen is described as uniformly pale brown. The size is not stated, nor the structure of the ungues.

Taeniorhynchus africanus, Neven-Lemaire, Arch. Parasit. 10, p. 271 (1906). This eannot be a true Taeniorhynchus, as the female has toothed claws.

Boycia mimomyiaformis. Newst., *l.c.* p. 34. Should probably be included in the genus *Ludlowia*; seems to be distinct from *L. pincerna*, which it most closely resembles, since the last *three* joints of the hind tarsi are described as *bronzy-ochrcous*, the legs otherwise being unbanded.

Neomelanoconion palpale, Newst., l.c. p. 31. This species clearly does not belong to the genus Neomelanoconion, which is a synonym of Culiciomyia. The palpi of the male are figured as two-jointed; if this is really the case, the species cannot be a Melanoconion. The figures of the male palpus and wing and the description of the antennae suggest a Ludlowia.

Index of Specific Names.

[Tabulated African species in heavy type : synonyms in italies.]

	P.	AGE.				P	AGE.
••• •••		251	annulipes, Wlk			•••	253
	256	, 261	annulirostris, Skuse.				256
		265	aquilus, Graham		•••		264
onia)		254	arabica, Giles (Mansor	nia)			249
dus)		246	arabicus, Becker (Culo	ex)			249
Taeniorhy	inchus)	266	aurites, Theo		•••		252
	256,	, 259	azoriensis, Theo				263
••• ••		247	bicolor, Mg			256,	26 6
	. 244.	, 245	bifoliata, Theo			•••	261
		247	bostocki, Theo				264
		259	caliginosus, Graham			246, 248,	250
		246	chamberlaini, Ludl.				244
		259	chloroventer, Theo.				264
	259,	,260	chrysothorax, Theo.				245
	252,	253	cinereus, Theo	•••		254,	255
	244.	251	clavipalpus, Theo.				244
		253	commorens, Walk.				246
	243, 256,	259	concolor, RD			261,	262
	 onia) dus) Taeniorhy 	P 256 256 dns) Taeniorhynchus) 256 244 244 259 259 259 259 252 244 244	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	PAGE. 251 annulipes, Wlk annulipostris. Skuse. annulirostris. Skuse. aquilus, Graham anabilitation 254 arabica, Giles (Mansoidans) 266 das) 246 rabica, Giles (Mansoidans) 266 aurites, Theo. 256, 259 256, 259 256, 259 256, 259 247 bicolor, Mg. 247 bicolor, Mg. 247 bostocki, Theo. 247 bostocki, Theo. 259 chloroventer, Theo. 259, 260 chrysothoras, Theo. 252, 253 cinerens, Theo. 252, 253 connerens, Walk. 244, 251 clavipalpus, Theo. 243, 256, 259 concolor, RD.	PAGE. 251 annulipes, Wlk 256, 261 annulirostris, Skuse. 266, 261 annulirostris, Skuse. 266 arabica, Giles (Mansonia) analis 244 arabica, Giles (Mansonia) dns) 246 arabica, Becker (Culex) Taeniorhynchus) 266 aurites, Theo. 256, 259 azoriensis, Theo. 247 bicolor, Mg. 247 bostocki, Theo. 247 bostocki, Theo. 259 chloroventer, Theo. 259 chloroventer, Theo. 252, 253 cinereus, Theo. 253	PAGE. 251 annulipes, Wlk 256, 261 annulirostris, Skuse. 265 aquilus, Graham mia) 256, 261 arabica, Giles (Mansonia) onia) 246 arabica, Giles (Mansonia) dns) 246 arabica, Becker (Culex) Taeniorhynchus) 266 aurites, Theo. 256, 259 azoriensis, Theo. 247 bicolor, Mg. 244 245 bifoliata, Theo. 247 bostocki, Theo. 247 bostocki, Theo. 259 chamberlaini, Ludl. 259 chloroventer, Theo. .	PAGE. P. 251 annulipes, Wlk 256 , 261 annulirostris, Skuse. 256 , 261 annulirostris, Skuse. 265 $aquilus$, Graham $aquilus$, Graham dns) 244 arabicus, Becker (Culex) Taeniorbynchus) 266 aurites, Theo. Taeniorbynchus) 266 aurites, Theo. <

266

			Т	AGE	1	PAGE
condulader Grünh			950	960	metallicus Theo 943 97	9 956
aongimilig Nowst		• • •	912 956	200	microannalatus Theo	961
constanting, incust.	•••	•••	440, 200	059	miniominiatus, raco,	4 966
cristatus, Theo	• • •	•••	202	, 400	minutus Theo	969
cumminsi, i neo.	•••	•••	Z47	, 240	Manutus, Theo	200
dalzieli, Theo	•••	••••	250	, 200		262
decens, Theo	•••	•••	257	,263	morsitans, Theo	251
dentatus, Theo	•••	•••	247	, 248	muciaus, Karsch	246
didieri, Neveu-Lemaire		•••		266	mundulus, Grünb	266
dissimilis, Theo		• • •	259	,260	neuvci, Theo	263
dorsalis, Mg			247	,248	nebulosa , Theo 254, 255, 26	3,266
drymoecius, Speiser.				253	nemorosus, Mg 24	7,248
durbanensis, Theo.			248	,250	neotaeniorhynchus, Theo	-261
duttoni, Theo			257, 259	,262	niger, Giles	247
euclastus, Theo				264	nigeriensis, Theo 24	8,250
fasciata, F			248	262	nigerrima. Theo	254
fasciolatus, Arrib.				252	nigra, Theo. (Mansonia) 25	3 254
fatigans Wied		•••	•••	969	wigra Theo (Chrysoconops)	959
freetowners Theo	•••		 954	955	nigrithoray Theo	959
fulyus Wind	•••		<i>40</i> 4	, 200 ara	nighter atas Theo 259.95	1 956
fuctors, will an	•••	•••	***	292	migrochaetalie, Theo 202, 20	4, 200
fuscopatpatis, 1 neo.	•••	•••		248	<i>nigrocostatis</i> , Theo	203
iuscopennatus, Theo.	•••	•••	252	,253	nocturnus, 1 neo	250
fuscum, Theo. (Proton	nelanoc	eoni	on) 244	,264	ocellata, Theo	250
fuscus, Theo. (Pectinop	alpus)	•••	•••	255	ochraceus, Theo. (Ochlerotatus) 246, 24	8,250
grahami, Theo. (Cule	x)	• • •	244, 257	,264	ochraceus, Theo. (Chrysoconops)	253
grahami, Theo. (Mucid	us)			246	ornatothoracis, Theo 257, 26	3,264
guiarti, Blanch	•••		257	,263	ornatus, Mg	-250
hirsutus, Theo.	7 • •		247	, 249	pallidocephala, Theo	-262
hirsutipalpis, Theo.			259	260	pallidostriatus, Theo	-250
hispida, Theo			244	245	pallidothoracis, Theo	263
inconspicuosa, Theo.				255	pallipes, Meg.	265
insignis, Carter			258	265	palpale. Newst	266
invenustus. Theo.			254	256	par Newst	266
invidiosus. Theo		•••	957	964	puruscelos Theo	250
jamaicensis Theo	•••	•••	201	050	nettiavavi Theo	1 969
kingii Theo	•••	•••	247	200	petityrew, Theo 25	9, 202 9, 902
latorolia Ma	•••	•••		200	pinterus, meo 250	5, 400 1 915
lauronti Nowat	•••	•••	248,	250	pinicerna, Granam 249	1, 240
laurently, Newst	•••	•••	***	266	pipiens, $11. \dots 241, 248, 251, 261$	1, 203
lencauthuise, LW.	•••	••••	***	249	plumosa, Theo 24-	, 245
leucarthrius, Speiser	•••	•••	248,	251	poicilipes, Theo	258
lineatopennis, Ludl.	•••	• • •	•••	245	pruina, Theo 257, 263	5,264
lividocostalis, Graham	* * *	•••	• > 0	263	pruinosns, Theo	263
longefurcatus, Beeker	•••	•••	•••	265	pseudoannulioris, Theo	259
longisquamosus, Theo).	•••	247,	249	pseudocinereus, Theo 254	1,255
lowisii, Theo	•••			250	pseudoconopas, Theo	253
luteolateralis, Theo.	•••			245	pulcripalpis, Rond 248	3, 249
maculicrura, Theo.				261	pulcritarsis, Rond 247	,249
maculiventris, Meq.				265	pullatus, Graham	264
maeulosa, Theo				248	punctings Theo	258
madagascariensis, Vent	•.		• • •	959	punctopes, theo	200
major. Theo.		••••	•••	200	punctocostans, 1 neo	245
mariac. Serg		•••	***	204	pusinus, meq	200
musenlus Theo	•••	•••	•••	249	pygmaens, Neven-Lemaire	266
medialineata Theo	•••	•••		203	quasigelidus, Theo. 243, 256, 258, 260	,264
modionmetata, 1 neo.	• • •	•••	253,	254	quasiguiarti, Theo	264
meanopanetata, 1 fieo,	•••	•••	•••	248	quasimodesta, Theo 256	, 266
meranorninus, Giles	• • •		•••	265	quasiunivittatus, Theo 248	,250

				$\mathbf{P}_{\mathbf{A}}$	AGE.				\mathbf{P}	AGE.
rima, Theo.	•••		•••	255, 258,	265	taeniorhynchoides, Gil	es	•••		258
rubinotus, Theo).			258,	265	taeniorhynchus, Wied	(Ochle	erotatus	s) 5	247,
rufinus, Big.			•••		265				251,	252
salisburiensis,	Theo.			258,	264	taeniorhynchus, Arrib.	(Taen	iorhynd	ehus)	251
salsus, Theo.	•••				261	tenax, Theo			256,	259
salus, Theo.					261	thalassius, Theo.	•••	•••	257,	261
sarawaki, Theo.	•••				259	theileri, Theo	•••	250, 2	57, 260,	262
scapularis, Rond.	•••			247,	256	tigripes, Grandpré		243, 2	51, 257,	261
scatophagoides.	Theo				246	titillans, Wlk	•••		251,	252
sergenti, Theo.	•••			258,	265	transraalensis, Theo.		•••		249
simplex, Theo.	•••			262,	263	uniformis, Theo. (Cul	liciomy	via)		255
simpsoni, Theo.				257,	263	uniformis, Theo. (M	ansoni	a)	253,	254
sinensis, Theo.					259	univittatus, Theo.		28	57, 260,	262
somaliensis, Ne	veu-Le	maire		257,	261	varioannulatus, Theo.			•••	263
spathipalpis, R	ond.			242, 243,	251	vigilax, Skuse	•••		247,	250
stoehri, Theo.					263	violaceus, Theo				252
subtilis, Serg.					248	viridis, Theo				263
sudanensis, Theo.			•••		246	willcocksi, Theo				248
taeniarostris, T	heo (I	Banksir	iell	a)	245	zeltneri, Neveu-Lema	ire			266
taeniarostris, The	o. (Gra	ıbhami	a)		259	zombaensis, Theo.			257,	264
						,				

Note on some Culicidae described by Dr. K. Grünberg.—In the Zoologischer Anzeiger for 1905 (pp. 377-390), Dr. K. Grünberg described six new *Culicidae* from Kamerun and Togo, and the following notes on them may be useful.

1. $Myzomyia \ bisignata$.—This is in all probability a synonym of M. rhodesiensis, Theo. The third costal spot seems fairly distinct in Grünberg's figure, while in the type of M. rhodesiensis it is quite small.

2. *M. unicolor.*—Without a doubt the same as *M. umbrosa*, Theo. In publishing my note on this species in the July number of this Bulletin, I overlooked the fact that *M. nili*, Theo. (First Rep. Welle, Lab., p. 66, 1904) is another synonym; the types correspond in every particular.

3. Myzorlynchus obscurus.—Evidently the same species which Theobald has subsequently described as *M. strachani*. As Grünberg suggests, it is very likely only a variety of *M. barbirostris*.

4. Stegomyia longipalpis.—Redescribed by Graham as S. pollinctor. Transferred by Theobald to Kingia.

5. Culex condylodesmus.—Probably only C. duttoni. See above (p. 260).

6. C. mundulus.-Perhaps a Culiciomyia. See above (p. 266).

F. W. EDWARDS.