



## Recognition of *Lasioconops* Theobald, 1903 and *Oculeomyia* Theobald, 1907 as separate subgenera of the genus *Culex* Linnaeus, 1758 (Diptera: Culicidae)

RALPH E. HARBACH

Department of Science, Natural History Museum, Cromwell Road, London SW7 5BD, UK.

✉ [r.harbach@nhm.ac.uk](mailto:r.harbach@nhm.ac.uk); <https://orcid.org/0000-0003-1384-6972>

In a series of studies on the pupal stage of mosquitoes that occur in Japan, Tanaka (2004) resurrected *Oculeomyia* Theobald, 1907 from synonymy with *Culex* Linnaeus, 1758 for *Cx. bitaeniorhynchus* Giles, 1901a and *Cx. sinensis* Theobald, 1903a, members of the “*Bitaeniorhynchus* Subgroup” of Sirivanakarn (1976). Based on descriptions and figures of larvae in the works of Hopkins (1952), Belkin (1962), Bram (1967) and Sirivanakarn (1976), Tanaka tacitly included 14 other species in the subgenus: *Cx. albinervis* Edwards, 1929; *Cx. annulioris* Theobald, 1901; *Cx. cornutus* Edwards, 1922; *Cx. epidesmus* (Theobald, 1910); *Cx. ethiopicus* Edwards, 1912 (synonymized with *bitaeniorhynchus* by Harbach 1988); *Cx. geminus* Colless, 1955; *Cx. infula* Theobald, 1901; *Cx. kinabaluensis* Sirivanakarn, 1976; *Cx. longicornis* Sirivanakarn, 1976; *Cx. luzonensis* Sirivanakarn, 1976; *Cx. pseudosinensis* Colless, 1955; *Cx. selangorensis* Sirivanakarn, 1976; *Cx. squamosus* (Taylor, 1914); and *Cx. starckeae* Stone & Knight, 1958. The *Bitaeniorhynchus* Subgroup, based on equivalence with the “*bitaeniorhynchus* series” of Edwards (1932), also includes the Afrotropical *Cx. aurantapex* Edwards, 1914, *Cx. giganteus* Ventrillon, 1906 and *Cx. poicilipes* (Theobald, 1903a), and the Oriental *Cx. samoensis* (Theobald, 1914). Tanaka did not specifically mention those four species except to say that “*Culex poicilipes* may belong to a different group.” In agreement with Tanaka, Harbach (1988) explicitly excluded *Cx. poicilipes* from the *Bitaeniorhynchus* Subgroup and provisionally included it in the *Sitiens* Subgroup (“*sitiens* subgroup” of Sirivanakarn 1976; “*sitiens* series” of Edwards 1932). There is no denying that taxonomic acts are sometimes overlooked or forgotten, and this is the vexatious reason why *Cx. poicilipes* was subsequently listed by Harbach (2011) as a member of the *Bitaeniorhynchus* Subgroup, and its recognition as a species of the subgenus *Oculeomyia* in the lexicon of Harbach (2018), the catalog of Wilkerson *et al.* (2021) and the online Mosquito Taxonomic Inventory (Harbach 2023). It is noteworthy that Edwards (1941) implied that *Cx. poicilipes* was not a member of the *Bitaeniorhynchus* Subgroup when he stated “*Culex bitaeniorhynchus* typifies a small group of species which are in most respects similar to *C. poicilipes* as described above, but lack the rows of small pale spots on the femora and tibiae”.

*Oculeomyia* was originally proposed as a distinct genus by Theobald (1907), with *Oculeomyia sarawaki* Theobald, 1907, a subjective junior synonym of *Culex infula* Theobald, 1901, as the type species (by monotypy). Edwards (1911) synonymized *Oculeomyia sarawaki* with *Taeniorhynchus ager* Giles, 1901b and later (Edwards 1913) synonymized both with *Cx. bitaeniorhynchus*, thus relegating *Oculeomyia* to synonymy with *Culex*. *Oculeomyia* remained in synonymy with *Culex sensu stricto* until Tanaka (2004) resurrected it from synonymy and validated it as a subgenus, with *Cx. infula* as the valid taxonomic type species. *Oculeomyia* was subsequently understood to include *Cx. poicilipes* and the other 18 species listed above.

Berlov & Kuberskaya (2023) recently published a paper in which they pointed out errors in the Taxonomic Catalog of Culicidae, Part Three of the *Mosquitoes of the World* (Wilkerson *et al.* 2021, Volume 2). The authors noted that *Cx. poicilipes* (Theobald, 1903a) is the type species of the nominal genus *Lasioconops* Theobald, 1903b, which was published earlier than *Oculeomyia* Theobald, 1907, and therefore replaced *Oculeomyia*. When Tanaka (2004) resurrected *Oculeomyia* from synonymy with *Culex* and validated it as a subgenus for species of the *Bitaeniorhynchus* Subgroup, he cited Harbach (1988) in his discussion of the pupa of *Cx. bitaeniorhynchus* but he apparently overlooked the fact that Harbach had removed *Cx. poicilipes* from the *Bitaeniorhynchus* Subgroup, otherwise he probably would have explicitly excluded it from *Oculeomyia*. Since *Cx. poicilipes* was not included as a member of the *Bitaeniorhynchus* Subgroup when Tanaka raised it to subgeneric status, the subgeneric name *Oculeomyia* should not have been replaced by *Lasioconops*. Harbach (1988) removed *Cx. poicilipes* from the *Bitaeniorhynchus* Subgroup and provisionally included it in the *Sitiens* Subgroup (Bram 1967) based on morphological similarities of the adults, male genitalia and larvae with the aberrant *Culex whitmorei* (Giles, 1904).

Tanaka (2004) noted that “the subgenus *Oculeomyia* is best characterized in the larval stage” and emphasized characteristics of the mandibles and maxillae of *Cx. bitaeniorhynchus* and *Cx. sinensis* observed by Tanaka *et al.* (1979) in his description of the subgenus. Since the mouthparts of other species of the subgenus had not been studied, Tanaka’s implicit inclusion of the species noted above, and his reluctance to include *Cx. poicilipes*, was based on “the size of the head, fused labrum [median labral plate], peculiar mentum plate [dorsomentum] and pecten”. As the information in Table 1, extracted from the descriptions and illustrations of Barraud (1934), Edwards (1941), Hopkins (1952), Belkin (1962), Brunhes (1967), Sirivanakarn (1976) and Harbach (1988), demonstrably shows, *Cx. poicilipes* is very dissimilar to the other 18 species included in *Oculeomyia*. Additionally, it is noteworthy that the larva of *Cx. poicilipes* exhibits an unusual behaviour associated with its modified siphon. Pronounced forward curvature of the siphon is associated with feeding from an inverted position, whereby the siphon points forward toward the head and the curvature may help to prevent damage to the spiracular apparatus while the long posterior elements of seta 1-S prevent the siphon from coming into contact with the substrate (Lewis 1949). The lobes of the spiracular apparatus are exceptionally large and apparently allow the larva to obtain oxygen from bubbles on the surface of submerged plant parts (Hopkins 1952).

**TABLE 1.** Striking morphological differences between *Cx. poicilipes* and the other species currently placed in the subgenus *Oculeomyia*.

Life stage	Characteristics	<i>Cx. poicilipes</i>	Species of <i>Oculeomyia</i> (excluding <i>Cx. poicilipes</i> )
Larvae	Clypeolabral suture	Present, median labral plate distinctly separated from dorsal apotome	Absent, median labral plate indistinguishably fused with dorsal apotome
	Dorsomentum	Broadly triangular with relatively few distinct teeth	In the shape of an equilateral or isosceles triangle with numerous minute teeth on each side
	Setae 4–6-C	Inserted posterior to mid-length of head capsule, relatively distant from seta 7-C	Inserted anterior to mid-length of head capsule, relatively close to seta 7-C
	Seta 2-VI	Inserted lateral to seta 3-VI	Inserted mesad of seta 3-VI
	Seta 1-III–VIII	Strongly developed, multi-branched and distinctly longer than half length of segment	Weakly developed, with few branches and much shorter than half length of segment
	Siphon	Short, stout, with pronounced forward curvature, index 3.5–4.7 (mean 4.2)	Long, slender, more or less straight, index 5.4–20.0 (mean 9.3)
	Pecten	8-14 spines (mean 11) borne on basal 0.33 of siphon	3–12 spines (mean 4 or 5) borne within basal 0.25 of siphon, usually < 0.15
	Seta 1-S	5 setae, 4 inserted at posterior midline, considerably longer than diameter of siphon, fifth seta inserted far lateral to 2 middle posterior setae	3-5 setae inserted in posterolateral row (penultimate seta may be slightly out of line), no longer than diameter of siphon, usually shorter
	Spiracular lobes	Unusually large	Normal, not unusually large
	Anal papillae	Either dorsal or ventral pair shorter than the other pair, longer pair no longer than saddle	Dorsal and ventral pairs same length, as long as or longer than saddle, usually longer
Pupae	Trumpet	More or less cylindrical, pinna short, 0.1–0.2 length of trumpet	Strongly flared, funnel-shaped, pinna long, about 0.4 length of trumpet
	Setae 6-V,VI	With 3–6 branches	Single
Adults	Legs	Anterior surfaces of femora and tibiae with row of pale spots	Anterior surfaces of femora and tibia entirely dark-scaled or more often with speckling of pale scales

The results of the phylogenetic study of Harbach *et al.* (2012) provide further evidence that *Cx. poicilipes* is distinct from the other members of the subgenus *Oculeomyia*. The authors explored the relationships among the genera and

subgenera of the tribe Culicini using 169 morphological characters scored for 86 species, including *Cx. bitaeniorhynchus*, *Cx. infula* and *Cx. poicilipes*. Thirteen “generally well-supported and stable” clades were recovered in three separate analyses. One of the clades included the three species in a sister relationship to species of the Mimeticus Subgroup (“mimeticus series” of Edwards 1932), with *Cx. poicilipes* as the sister of *Cx. bitaeniorhynchus* + *Cx. infula*.

As a result of the recent revisionary study of Harbach & Wilkerson (2023), the subgenus *Oculeomyia* now also includes *Cx. consimilis* Newstead, 1907 (in Newstead *et al.* 1907), *Cx. ellinorae* Ovazza, Hamon & Neri, 1956, *Cx. jinjaensis* Edwards, 1941 and *Cx. major* Edwards, 1935. *Culex consimilis* was previously recognized as a subspecies of *Cx. annulioris*, *Cx. ellinorae* and *Cx. jinjaensis* as subspecies of *Cx. aurantapex*, and *Cx. major* as a synonym of *Cx. consimilis*.

Based on information provided above and the morphological distinctions listed in Table 1, it is obvious that the 23 species presently included in the subgenus *Oculeomyia* belong to two separate phyletic lines. Since it appears that *Cx. poicilipes* is unrelated to the other species presently placed in *Oculeomyia*, as well as to species of the subgenus *Culex*, *Cx. poicilipes* is hereby formally recognized as the monotypic member of a distinct subgenus: *Lasioconops* Theobald, 1903b **stat. nov.**, denoted by the three-letter abbreviation *Lsi*. The genus- and species-group names included in the subgenera *Lasioconops* and *Oculeomyia* as a result of this action are summarized in the following lists.

### ***Lasioconops* Theobald, 1903b**

TYPE SPECIES: *Lasioconops poicilipes* Theobald, 1903b.

SYNONYMS:

*Pseudoheptaphlebomyia* Ventrillon, 1905 (type species: *P. madagascariensis* Ventrillon, 1905; synonymy with *Culex quasigelidus* Theobald, 1903a by Edwards 1911; synonymy of *Cx. quasigelidus* with *Cx. poicilipes* by Edwards 1932).

*Aporoculex* Theobald, 1907 (type species: *A. punctipes* Theobald, 1907; synonymy with *Cx. quasigelidus* by Edwards 1911; synonymy of *Cx. quasigelidus* with *Cx. poicilipes* by Edwards 1932).

INCLUDED SPECIES: *Culex poicilipes* (Theobald, 1903b).

### ***Oculeomyia* Theobald, 1907**

TYPE SPECIES: *Oculeomyia sarawaki* Theobald, 1907 (synonymy with *Culex infula* Theobald, 1901 by Sirivanakarn 1976).

SYNONYM:

*Trichopronomyia* Theobald, 1905 (type species: *T. annulata* Theobald, 1905; synonymy with *Culex squamosus* (Taylor, 1914) by Edwards 1924).

INCLUDED SPECIES: *Culex albinervis* Edwards, 1929; *Cx. annulioris* Theobald, 1901; *Cx. aurantapex* Edwards, 1914; *Cx. bitaeniorhynchus* Giles, 1901a; *Cx. consimilis* Newstead, 1907; *Cx. cornutus* Edwards, 1922; *Cx. ellinorae* Ovazza, Hamon & Neri, 1956; *Cx. epidesmus* (Theobald, 1910); *Cx. geminus* Colless, 1955; *Cx. giganteus* Ventrillon, 1906; *Cx. infula* Theobald, 1901; *Cx. kinabaluensis* Sirivanakarn, 1976; *Cx. longicornis* Sirivanakarn, 1976; *Cx. luzonensis* Sirivanakarn, 1976; *Cx. major* Edwards, 1935; *Cx. pseudosinensis* Colless, 1955; *Cx. samoensis* (Theobald, 1914); *Cx. selangorensis* Sirivanakarn, 1976; *Cx. sinensis* Theobald, 1903a; *Cx. squamosus* (Taylor, 1914); *Cx. starckeae* Stone & Knight, 1958.

With the exclusion of *Cx. (Lsi.) poicilipes* (Theobald, 1903a), the subgenus *Oculeomyia* now includes 22 species. However, it should be noted that Lee *et al.* (1989), without explanation, listed *Cx. samoensis* as a species of the Sitiens Group without placement in a subgroup. Despite being known from only three adult females (one lost), Belkin (1962) noted that “*C. samoensis* seems to be a distinct species” but it was “possible that *samoensis* is the Samoan representative of *albinervis*, which it closely resembles.” However, until proven otherwise, *Cx. samoensis* should continue to be classified as a species of the subgenus *Oculeomyia*.

### **Acknowledgement**

I gratefully acknowledge Neal Evenhuis (Bishop Museum, Honolulu) and an anonymous reviewer for their comments and suggestions for improving to the manuscript. Anders Lindström, National Veterinary Institute, Uppsala, Sweden kindly provided the cover image of *Culex pipiens* displayed with this publication on the Zootaxa webpage.

### **References**

Barraud, P.J. (1934) *The fauna of British India, including Ceylon and Burma. Diptera. Vol. V. Family Culicidae. Tribes Megarhinini and Culicini.* Taylor and Francis, London, xxviii + 463 pp., 8 pls.

- Belkin, J.N. (1962) *The mosquitoes of the South Pacific (Diptera, Culicidae)*. Vols 1 & 2. University of California Press, Berkeley and Los Angeles, California, xii + 608 pp., 412 figs.
- Berlov, O.E. & Kuberskaya, O.V. (2023) Additions and corrections to the Catalog of blood-sucking mosquitoes of the world (Insecta: Diptera, Culicidae) by Wilkerson et al. (2021). *Amurian Zoological Journal*, 15 (1), 110–118. [in Russian] <https://doi.org/10.33910/2686-9519-2023-15-1-110-118>
- Bram, R.A. (1967) Contributions to the mosquito fauna of Southeast Asia. II. The genus *Culex* in Thailand (Diptera: Culicidae). *Contributions of the American Entomological Institute*, 2 (1), 1–296.
- Brunhes, J. (1967) Contribution a l'étude des culicides de Madagascar. Description des larves et nymphes de *Culex (C.) giganteus* Ventrillon et *Culex (C.) argenteopunctatus* Ventrillon [sic]. Notes complémentaires sur les adultes de ces deux espèces. *Cahiers ORSTOM Série Entomologie médicale et Parasitologie*, 5 (4), 207–216.
- Colless, D.H. (1955) Notes on the culicine mosquitoes of Singapore. I.—Three new species of *Culex* (Diptera, Culicidae) and a redescription of *Culex hutchinsoni* Barraud. *Annals of Tropical Medicine and Parasitology*, 49 (3), 311–319. <https://doi.org/10.1080/00034983.1955.11685680>
- Edwards, F.W. (1911) The African species of *Culex* and allied genera. *Bulletin of Entomological Research*, 2 (3), 241–268. <https://doi.org/10.1017/S0007485300001450>
- Edwards, F.W. (1912) A synopsis of the species of African Culicidae, other than *Anopheles*. *Bulletin of Entomological Research*, 3 (1), 1–53. <https://doi.org/10.1017/S0007485300001656>
- Edwards, F.W. (1913) New synonymy in Oriental Culicidae. *Bulletin of Entomological Research*, 4 (3), 221–242. <https://doi.org/10.1017/S00074853000043133>
- Edwards, F.W. (1914) New species of Culicidae in the British Museum, with notes on the genitalia of some African *Culex*. *Bulletin of Entomological Research*, 5 (1), 63–81. <https://doi.org/10.1017/S0007485300034568>
- Edwards, F.W. (1922) A synopsis of adult Oriental culicine (including megarhinine and sabethine) mosquitoes. Part I. *Indian Journal of Medical Research*, 10 (1), 249–293, 3 pls.
- Edwards, F.W. (1924) A synopsis of the adult mosquitos of the Australasian Region. *Bulletin of Entomological Research*, 14 (4), 351–401. <https://doi.org/10.1017/S0007485300045624>
- Edwards, F.W. (1929b) Mosquito notes.—VIII. *Bulletin of Entomological Research*, 20 (3), 321–343. <https://doi.org/10.1017/S0007485300021283>
- Edwards, F.W. (1932) *Genera Insectorum. Diptera, Fam. Culicidae. Fascicle 194*. V. Verteneuil & L. Desmet, Bruxelles [Brussels], 258 pp., 5 pls.
- Edwards, F.W. (1935) Mosquito notes.—XII. *Bulletin of Entomological Research*, 26 (2), 127–136. <https://doi.org/10.1017/S0007485300038116>
- Edwards, F.W. (1941) *Mosquitoes of the Ethiopian Region III.—Culicine adults and pupae*. Printed by order of the Trustees, British Museum (Natural History), London, viii + 499 pp.
- Giles, G.M. (1901a) A plea for the collective investigation of Indian Culicidae, with suggestions as to moot points for enquiry, and a prodromus of species known to the author. *Journal of the Bombay Natural History Society*, 13, 592–610, 2 pls.
- Giles, G.M. (1901b) Six new species of Culicidae from India. *The Entomologist*, 34 (458), 192–197.
- Giles, G.M. (1904) Notes on some collections of mosquitoes, &c., received from the Philippine Islands and Angola; with some incidental remarks upon classification. *Journal of Tropical Medicine*, 7 (28), 365–369.
- Harbach, R.E. (1988) The mosquitoes of the subgenus *Culex* in southwestern Asia and Egypt (Diptera: Culicidae). *Contributions of the American Entomological Institute*, 24 (1), i–vi + 1–240.
- Harbach, R.E. (2011) Classification within the cosmopolitan genus *Culex* (Diptera: Culicidae): The foundation for molecular systematics and phylogenetic research. *Acta Tropica*, 120 (1–2), 1–14. <https://doi.org/10.1016/j.actatropica.2011.06.005>
- Harbach, R.E. (2018) *Culicipedia: species-group, genus-group and family-group names in Culicidae (Diptera)*. CABI, Wallingford, Oxfordshire, xviii + 378 pp. <https://doi.org/10.1079/9781786399052.0000>
- Harbach, R.E. (2023) Subgenus *Oculeomyia* Theobald, 1907. Mosquito Taxonomic Inventory. Available from: <https://mosquito-taxonomic-inventory.myspecies.info/simpletaxonomy/term/6186> (accessed 24 April 2023)
- Harbach, R.E. & Wilkerson, R.C. (2023) The insupportable validity of mosquito subspecies (Diptera: Culicidae) and their exclusion from culicid classification. *Zootaxa*, 5303 (1), 1–184. <https://doi.org/10.11646/zootaxa.5303.1.1>
- Harbach, R.E., Kitching, I.J., Culverwell, C.L., Dubois, J. & Linton, Y.-M. (2012) Phylogeny of mosquitoes of tribe Culicini (Diptera: Culicidae) based on morphological diversity. *Zoologica Scripta*, 41 (5), 499–514. <https://doi.org/10.1111/j.1463-6409.2012.00546.x>
- Hopkins, G.H.E. (1952) *Mosquitoes of the Ethiopian Region I.—Larval bionomics of mosquitoes and taxonomy of culicine larvae*. 2<sup>nd</sup> Edition. British Museum (Natural History), London, viii + 355 pp.
- Lee, D.J., Hicks, M.M., Debenham, M.L., Griffiths, M., Marks, E.N., Bryan, J.H. & Russell, R.C. (1989) *The Culicidae of the Australasian Region. Vol. 7. Nomenclature, synonymy, literature, distribution, biology and relation to disease. Genus Culex*.

- Subgenera Acallyntrum, Culex*. Monograph Series. entomology Monograph No. 2. Australian Government Publishing Service, Canberra, ix + 281 pp.
- Lewis, D.J. (1949) Tracheal gills in some African culicine mosquito larvae. *Proceedings of the Royal Entomological Society of London. Series A, General Entomology*, 24 (4–6), 60–66.  
<https://doi.org/10.1111/j.1365-3032.1949.tb00718.x>
- Linnaeus, C. (1758) *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Tomus 1. Editio decima. Laurentii Salvii, Holmiae [Stockholm], 824 pp.  
<https://doi.org/10.5962/bhl.title.542>
- Newstead, R., Dutton, J.E. & Todd, J.L. (1907) Insects and other Arthropoda collected in the Congo Free State. *Annals of Tropical Medicine and Parasitology*, 1 (1–5), 3–113.  
<https://doi.org/10.1080/00034983.1907.11719252>
- Ovazza, M., Hamon, J. & Neri, P. (1956) Contribution a l'étude des diptères vulnérants de l'empire d'Éthiopie. I. Culicidae. *Bulletin de la Société de Pathologie exotique*, 49 (1), 151–182.
- Sirivanakarn, S. (1976) Medical entomology studies—III. A revision of the subgenus *Culex* in the Oriental Region (Diptera: Culicidae). *Contributions of the American Entomological Institute*, 12 (2), i–iii + 1–272.
- Stone, A. & Knight, K.L. (1958) Two new names in mosquitoes (Diptera, Culicidae). *Proceedings of the Entomological Society of Washington*, 60 (2), 69.
- Tanaka, K. (2004) Studies on the pupal mosquitoes of Japan (10) *Culex (Culex)* (Diptera, Culicidae). *Japanese Journal of Systematics Entomology*, 10 (1), 9–42.
- Tanaka, K., Mizusawa, K. & Saugstad, E.S. (1979) A revision of the adult and larval mosquitoes of Japan (including the Ryukyu Archipelago and the Ogasawara Islands) and Korea (Diptera: Culicidae). *Contributions of the American Entomological Institute*, 16, i–vii + 1–987.
- Taylor, F.H. (1914) The *Culicidae [sic]* of Australia.—I. *Transactions of the Entomological Society of London*, 1913, 683–708, 4 pls.  
<https://doi.org/10.1111/j.1365-2311.1914.tb01794.x>
- Theobald, F.V. (1901) *A monograph of the Culicidae or mosquitoes*. Vol. 1. British Museum (Natural History), London, xviii + 424 pp., viii pls., pls I–XXXVII and A–E under separate cover.  
<https://doi.org/10.5962/bhl.title.58067>
- Theobald, F.V. (1903a) *A monograph of the Culicidae or mosquitoes*. Vol. 3. British Museum (Natural History), London, xvii + 1 (errata) + 359 pp., foldout table, 17 pls.  
<https://doi.org/10.5962/bhl.title.58067>
- Theobald, F.V. (1903b) Report on a collection of mosquitoes or Culicidae, etc, from Gambia, and descriptions of new species. *Memoirs of the Liverpool School of Tropical Medicine*, 10 (Appendix), i–xi, 2 pls.
- Theobald, F.V. (1905) A catalogue of the Culicidae in the Hungarian National Museum, with descriptions of new genera and species. *Annales Musei Nationalis Hungarici*, 3 (1), 61–119, figure legends, 4 pls.
- Theobald, F.V. (1907) *A monograph of the Culicidae or mosquitoes*. Vol. 4. British Museum (Natural History), London, xix + 639 pp., 16 pls.  
<https://doi.org/10.5962/bhl.title.58067>
- Theobald, F.V. (1910) Second report on the collection of Culicidae in the Indian Museum, Calcutta, with descriptions of new genera and species. *Records of the Indian Museum*, 4 (1), 1–33, 3 pls.
- Theobald, F.V. (1914) A new mosquito from Samoa. *The Entomologist*, 47 (608), 36–37.
- Ventrillon, E. (1905) Culicidés nouveaux de Madagascar. *Bulletin du Muséum National d'Histoire Naturelle*, 11 (6), 427–431.
- Ventrillon, E. (1906) *Culex* nouveaux de Madagascar. *Bulletin du Muséum National d'Histoire Naturelle*, 12 (2), 100–106.
- Wilkerson, R.C., Linton, Y.-M. & Strickman, D. (2021) *Mosquitoes of the world*. Vols. 1 & 2. Johns Hopkins University Press, Baltimore, Maryland, 1332 pp.  
<https://doi.org/10.1186/s13071-021-04848-6>